arXiv:1112.2399v2 [math.RT] 1 Jul 2012

NILPOTENT ORBITS IN SMALL CHARACTERISTIC

TING XUE

Abstract. We show that the number of nilpotent orbits in the dual of an exceptional Lie algebra is finite in bad characteristic. We determine the closure relations on the set of nilpotent orbits in the dual of classical and exceptional Lie algebras. Moreover for classical groups, we give an explicit description of the nilpotent pieces (which are unions of nilpotent orbits) in the dual defined in [L5, X4], in particular the definition of nilpotent pieces in [L5, X4] coincides with the definition given by closure relations on nilpotent orbits.

1. INTRODUCTION

Let G be a connected reductive algebraic group defined over an algebraically closed field $\mathbf{k}$ of characteristic $p \geq 0$. Let $\mathfrak{g}$ be the Lie algebra of G and $\mathfrak{g}^*$ the dual vector space of $\mathfrak{g}$. Denote by $\mathcal{N}_{\mathfrak{g}^*}$ the set of nilpotent elements in $\mathfrak{g}^*$ (recall that an element $\xi : \mathfrak{g} \to \mathbf{k}^*$ is called nilpotent if it annihilates some Borel subalgebra of $\mathfrak{g}$, see [KW]). Note that G acts on $\mathcal{N}_{\mathfrak{g}^*}$ by coadjoint action. It is known that the number of G-orbits in $\mathcal{N}_{\mathfrak{g}^*}$ is finite when p is a good prime for G (in this case we can identify $\mathfrak{g}^*$ with $\mathfrak{g}$ via a G-invariant non-degenerate bilinear form on $\mathfrak{g}$ and the corresponding result for $\mathfrak{g}$ is well-known) or when G is of type B, C or D ([X1]). It remains to study the number of G-orbits in $\mathcal{N}_{\mathfrak{g}^*}$ when G is of exceptional type and p is a bad prime for G. One of our results is that this number is again finite. (This result has appeared in the PhD thesis of the author. We include it here for completeness.) Let $\underline{\mathcal{N}_{\mathfrak{g}^*}}$ denote the set of G-orbits in $\mathcal{N}_{\mathfrak{g}^*}$ (nilpotent orbits in $\mathfrak{g}^*$). On $\underline{\mathcal{N}_{\mathfrak{g}^*}}$ there are natural order relations; for two orbits c and c', we say that $\mathrm{c} \leq \mathrm{c}'$ if c is contained in the closure of c', and that $\mathrm{c} < \mathrm{c}'$ if $\mathrm{c} \leq \mathrm{c}'$ and $\mathrm{c} \neq \mathrm{c}'$. We determine the order relations on $\underline{\mathcal{N}_{\mathfrak{g}^*}}$ (when p is bad).

In [L5, X4] Lusztig and the author define a partition of $\mathcal{N}_{\mathfrak{g}^*}$ into nilpotent pieces (for classical groups), indexed by the set $\underline{\mathcal{U}_{G_{\mathbb{C}}}}$ of unipotent orbits in the group $G_{\mathbb{C}}$ over complex numbers of the same type as G. On the other hand, there is a natural injective map from the set $\underline{\mathcal{U}_{G_{\mathbb{C}}}}$ to the set $\underline{\mathcal{N}_{\mathfrak{g}^*}}$ (given by Springer correspondence). Using this map and the order relations on $\underline{\mathcal{N}_{\mathfrak{g}^*}}$ one can define a partition of $\mathcal{N}_{\mathfrak{g}^*}$ into locally closed pieces; following Lusztig we call them MS-pieces (after Mizuno and Spaltenstein). We show that for classical groups the MS-pieces are the same as nilpotent pieces defined by Lusztig and the author. In particular we determine which nilpotent orbits lie in the same piece (Proposition 6.1).

This paper is organized as follows. Sections 2-6 study the cases when G is a classical group and $p = 2$. In Section 2 we recall natural order relations on the set of irreducible characters of the Weyl group of G (which have been used by Spaltenstein [S2] to describe closure relations of nilpotent orbits in $\mathfrak{g}$); the classification of nilpotent orbits in $\mathfrak{g}^*$; the combinatorial description of the Springer correspondence maps; and the definitions of nilpotent pieces in $\mathfrak{g}^*$ given in [L5, X4]. In section 3 and Section 4, we describe the Springer fibers at elements in $\mathcal{N}_{\mathfrak{g}^*}$ and induction for nilpotent orbits in $\mathfrak{g}^*$ (by an easy adaptation of [LS, S2]) respectively which give preparation for Section 5. In Section 5 we describe the order relations on $\underline{\mathcal{N}_{\mathfrak{g}^*}}$ which are given by natural order relations on the set of irreducible characters of the Weyl group of G via Springer correspondence. In Section 6 we describe the nilpotent pieces in $\mathfrak{g}^*$ explicitly and show that they coincide with MS-pieces. In Section 7 we study the nilpotent orbits in $\mathfrak{g}^*$ when G is of exceptional type and p is bad for G; in particular, we show that the set $\underline{\mathcal{N}_{\mathfrak{g}^*}}$ is finite and we determine the closure relations on $\underline{\mathcal{N}_{\mathfrak{g}^*}}$.

Acknowledgement I would like to thank George Lusztig for helpful discussions and for encouragement.

2. NOTATIONS AND RECOLLECTIONS

2.1. **Order relations on the set of irreducible characters of Wely groups of type** B, C, D**.** For a finite group H we denote by $H^\wedge$ the set of irreducible characters of H (over $\mathbb{C}$).

Let $\mathcal{P}(n)$ denote the set $\{\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots) \,|\, |\lambda| := \sum_{i \geq 1} \lambda_i = n\}$ of all partitions of an integer n. For a partition $\lambda \in \mathcal{P}(n)$ and each $j \geq 1$, we set

$$\lambda_j^* = |\{\lambda_i \,|\, \lambda_i \geq j|\} \text{ and } m_\lambda(j) = \lambda_j^* - \lambda_{j+1}^*.$$

Let $\mathcal{P}_2(n)$ denote the set $\{(\mu)(\nu) \,|\, |\mu| + |\nu| = n\}$ of pairs of partitions. If W is a Weyl group of type B_n or C_n (resp. D_n), then we can identify $W^\wedge$ with the set $\mathcal{P}_2(n)$ (resp. the set $\{(\mu)(\nu) \in \mathcal{P}_2(n) \,|\, \text{if } i \text{ is the smallest integer such that } \mu_i \neq \nu_i, \text{ then } \nu_i < \mu_i\}$ with each pair $(\mu)(\mu)$ counted twice) ([L1]). There are natural order relations on the set $\mathcal{P}_2(n)$ as follows. We say that $(\mu)(\nu) \leq (\mu')(\nu')$ if

$$\sum_{i \in [1,j]} (\mu_i + \nu_i) \leq \sum_{i \in [1,j]} (\mu'_i + \nu'_i) \text{ and } \sum_{i \in [1,j-1]} (\mu_i + \nu_i) + \mu_j \leq \sum_{i \in [1,j-1]} (\mu'_i + \nu'_i) + \mu'_j \text{ for all } j \geq 1;$$

and that $(\mu)(\nu) < (\mu')(\nu')$ if $(\mu)(\nu) \leq (\mu')(\nu')$ and $(\mu)(\nu) \neq (\mu')(\nu')$. This gives rise to order relations on $W^\wedge$ (in the case of type D_n, the two degenerate characters corresponding to a pair $(\mu)(\mu)$ are incomparable).

2.2. **Classification of nilpotent orbits in** $\mathfrak{g}^*$ **(type** B, C, D**).** Let V be a finite dimensional vector space over $\mathbf{k}$ equipped with a fixed non-degenerate symplecitc from $\langle , \rangle$ (resp. a fixed non-degenerate quadratic form Q with the associated bilinear form denoted by β). We can assume that $G = Sp(V, \langle , \rangle)$ (resp. $G = SO(V, Q)$), the subgroup of $GL(V)$ (resp. identity component of the subgroup $O(V, Q)$ of $GL(V)$) that preserves $\langle , \rangle$ (resp. Q). Then $\mathfrak{g} = \mathfrak{sp}(V, \langle , \rangle) = \{x \in \mathfrak{gl}(V) \,|\, \langle xv, w \rangle + \langle v, xw \rangle = 0 \ \forall \ v, w \in V\}$ (resp. $\mathfrak{g} = \mathfrak{o}(V, Q) = \{x \in \mathfrak{gl}(V) \,|\, \beta(xv, v) = 0 \ \forall \ v \in V, \ x|_{\mathrm{Rad(Q)}} = 0\}$).

Let $\mathfrak{Q}(V)$ (resp. $\mathfrak{S}(V)$) denote the vector space of all quadratic forms $V \to \mathbf{k}$ (resp. all symplectic forms $V \times V \to \mathbf{k}$). We have a vector space isomorphism (see [L5, X4])

(a) $\quad \mathfrak{sp}(V, \langle , \rangle)^* \xrightarrow{\sim} \mathfrak{Q}(V), \ \xi \mapsto \alpha_\xi, \ \alpha_\xi(v) = \langle v, Xv \rangle$

(resp. $\mathfrak{o}(V, Q)^* \xrightarrow{\sim} \mathfrak{S}(V), \ \xi \mapsto \beta_\xi, \ \beta_\xi(v, w) = \beta(Xv, w) - \beta(v, Xw)$),

where X is such that $\xi(x) = \mathrm{tr}(Xx)$ for all $x \in \mathfrak{sp}(V, \langle , \rangle)$ (resp. $\mathfrak{o}(V, Q)$).

Suppose that $G = Sp(V, \langle , \rangle)$ with $\dim V = 2n$ and $\xi \in \mathcal{N}_{\mathfrak{g}^*}$. Let α_ξ always denote the quadratic form corresponding to ξ under the isomorphism in (a) and let β_ξ always denote the bilinear form associated to α_ξ (given by $\beta_\xi(v, w) = \alpha_\xi(v + w) - \alpha_\xi(v) - \alpha_\xi(w)$). Let $T_\xi : V \to V$ be defined by $\langle T_\xi v, w \rangle = \beta_\xi(v, w)$ for all $v, w \in V$. Assume that $p = 2$. Then the G-orbit of ξ is characterized by a pair (λ, χ) as follows ([X1]):

(i) the partition $\lambda \in \mathcal{P}(2n)$ given by the sizes of Jordan blocks of T_ξ (we have $m_\lambda(i)$ even for all $i > 0$);

(ii) the map $\chi : \mathbb{N} \to \mathbb{N}$ given by $\chi(k) := \chi_{\alpha_\xi}(k) = \min\{l \,|\, T_\xi^k v = 0 \Rightarrow \alpha_\xi(T_\xi^l v) = 0 \ \forall\ v \in V\}$ (we have $\frac{\lambda_i - 1}{2} \leq \chi(\lambda_i) \leq \lambda_i$, $\chi(\lambda_i) \geq \chi(\lambda_{i+1})$ and $\lambda_i - \chi(\lambda_i) \geq \lambda_{i+1} - \chi(\lambda_{i+1})$ for all $i \geq 1$).

Suppose that $G = SO(V,Q)$ and $\xi \in \mathcal{N}_{\mathfrak{g}}^*$. Let β_ξ always denote the symplectic form corresponding to ξ under the isomorphism in (a).

If $p = 2$, $G = SO(V,Q)$ and $\dim V = 2n$, let $T_\xi : V \to V$ be defined by $\beta(T_\xi v, w) = \beta_\xi(v,w)$ for all $v, w \in V$. The $O(V,Q)$-orbit of ξ is characterized by a pair (λ, χ) as follows ([H]):

(i) the partition $\lambda \in \mathcal{P}(2n)$ given by the sizes of Jordan blocks of T_ξ (we have $m_\lambda(i)$ even for all $i > 0$);

(ii) the map $\chi : \mathbb{N} \to \mathbb{N}$ given by $\chi(k) := \chi_{T_\xi}(k) = \min\{l \,|\, T_\xi^k v = 0 \Rightarrow Q(T_\xi^l v) = 0 \ \forall\ v \in V\}$ (we have $\frac{\lambda_i}{2} \leq \chi(\lambda_i) \leq \lambda_i$, $\chi(\lambda_i) \geq \chi(\lambda_{i+1})$ and $\lambda_i - \chi(\lambda_i) \geq \lambda_{i+1} - \chi(\lambda_{i+1})$ for all $i \geq 1$).

Assume that $p = 2$, $G = SO(V,Q)$ and $\dim V = 2n+1$. Let $m \in [0,n]$ be the unique integer such that there exists a (unique) set of vectors $\{v_i, i \in [0,m]\}$ with

(b) $Q(v_m) = 1,\ Q(v_i) = 0,\ \beta_\xi(v_i, v) = \beta(v_{i-1}, v),\ i \in [1,m],\ \beta(v_m, v) = 0,\ \beta_\xi(v_0, v) = 0,\ \forall\ v \in V.$

If $m = 0$, let W be a complementary subspace of $\mathrm{span}\{v_0\}$ in V; if $m \geq 1$, let $\{u_i, i \in [0, m-1]\}$ be a set of vectors such that

(c) $Q(u_0) = 0, \beta(u_0, v_j) = \delta_{j,0}, j \in [0,m]; \beta_\xi(u_{i-1}, v) = \beta(u_i, v), Q(u_i) = 0, i \in [1, m-1], \forall\ v \in V$

and let $W = \{v \in V \,|\, \beta(v, u_i) = \beta(v, v_i) = \beta_\xi(u_0, v) = 0\}$. Then $V = \mathrm{span}\{u_i, v_i\} \oplus W$ and $\beta|_W$ is non-degenerate. Define $T_\xi : W \to W$ by $\beta_\xi(w, w') = \beta(T_\xi w, w')$ for all $w, w' \in W$ and let $\chi_W : \mathbb{N} \to \mathbb{N}$ be given by $\chi_W(k) = \min\{l \,|\, T_\xi^k w = 0 \Rightarrow Q(T_\xi^l w) = 0 \ \forall\ w \in W\}$. Then the orbit of ξ is characterized by $(m; (\lambda, \chi))$ as follows ([X1]):

(i) the integer $m \in [0,n]$;

(ii) the partition $\lambda \in \mathcal{P}(2n - 2m)$ given by the sizes of Jordan blocks of T_ξ (we have $m_\lambda(i)$ even for all $i > 0$);

(iii) the map $\chi : \mathbb{N} \to \mathbb{N}$ given by $\chi(i) = \max(i - m, \chi_W(i))$ (we have $m \geq \lambda_i - \chi(\lambda_i) \geq \lambda_{i+1} - \chi(\lambda_{i+1})$, $\frac{\lambda_i}{2} \leq \chi(\lambda_i) \leq \lambda_i$ and $\chi(\lambda_i) \geq \chi(\lambda_{i+1})$ for all $i \geq 1$).

Note that $(m; (\lambda, \chi))$ does not depend on the choice of W and u_0.

Let $\mathfrak{N}_{B_n}^{*2}$ (resp. $\mathfrak{N}_{C_n}^{*2}$, $\mathfrak{N}_{D_n}^{*2}$) be the set of all $(m; (\lambda, \chi))$ (resp. (λ, χ)) corresponding to nilpotent orbits ($p = 2$) in $\mathfrak{o}(2n+1)^*$ (resp. $\mathfrak{sp}(2n)^*$, $\mathfrak{o}(2n)^*$). Note that in the case of $\mathfrak{o}(2n)^*$, there are two orbits corresponding to each pair (λ, χ) with $\chi(\lambda_i) = \frac{\lambda_i}{2}$ for all i. Let

$$\mathfrak{N}_B^{*2} = \cup_{n \geq 0} \mathfrak{N}_{B_n}^{*2}, \quad \mathfrak{N}_C^{*2} = \cup_{n \geq 0} \mathfrak{N}_{C_n}^{*2}, \quad \mathfrak{N}_D^{*2} = \cup_{n \geq 0} \mathfrak{N}_{D_n}^{*2}.$$

2.3. **Combinatorial description of Springer correspondence maps (type B, C, D).** Let W denote the Weyl group of G. Recall that we have an injective Springer correspondence map [X1]:

$$\gamma_{\mathfrak{g}^*} : \underline{\mathcal{N}_{\mathfrak{g}^*}} \to W^\wedge,$$

which maps an orbit c to the Weyl group character corresponding to the pair $(\mathrm{c},1)$ under Springer correspondence. When $p=2$, the Springer correspondence maps $\gamma_{\mathfrak{g}^*}$ are given as follows ([X3])

$$\begin{aligned}\gamma^*_{B_n} &:= \gamma_{\mathfrak{o}(2n+1)^*}: \underline{\mathcal{N}_{\mathfrak{o}(2n+1)^*}} = \mathfrak{N}^{*2}_{B_n} \to W^\wedge,\\ &(m;(\lambda,\chi)) \mapsto (\mu)(\nu),\ \mu_1 = m,\ \mu_{i+1} = \lambda_{2i-1} - \chi(\lambda_{2i-1}),\ \nu_i = \chi(\lambda_{2i-1}),\ i \geq 1;\end{aligned}$$

$$\begin{aligned}\gamma^*_{C_n} &:= \gamma_{\mathfrak{sp}(2n)^*}: \underline{\mathcal{N}_{\mathfrak{sp}(2n)^*}} = \mathfrak{N}^{*2}_{C_n} \to W^\wedge,\\ &(\lambda,\chi) \mapsto (\mu)(\nu),\ \mu_i = \chi(\lambda_{2i-1}),\ \nu_i = \lambda_{2i-1} - \chi(\lambda_{2i-1}),\ i \geq 1;\\ \gamma^*_{D_n} &:= \gamma_{\mathfrak{o}(2n)^*}: \underline{\mathcal{N}_{\mathfrak{o}(2n)^*}} = \mathfrak{N}^{*2}_{D_n} \to W^\wedge,\\ &(\lambda,\chi) \mapsto (\mu)(\nu),\ \mu_i = \chi(\lambda_{2i-1}),\ \nu_i = \lambda_{2i-1} - \chi(\lambda_{2i-1}),\ i \geq 1.\end{aligned}$$

We denote the image of $\gamma^*_{B_n}$ (resp. $\gamma^*_{C_n}$, $\gamma^*_{D_n}$) (when $p=2$) by $\mathfrak{X}^{*2}_{B_n}$ (resp. $\mathfrak{X}^{*2}_{C_n}$ and $\mathfrak{X}^{*2}_{D_n}$). Let $\mathfrak{X}^{*2}_B = \cup_{n\geq 0}\mathfrak{X}^{*2}_{B_n}$, $\mathfrak{X}^{*2}_C = \cup_{n\geq 0}\mathfrak{X}^{*2}_{C_n}$ and $\mathfrak{X}^{*2}_D = \cup_{n\geq 0}\mathfrak{X}^{*2}_{D_n}$.

2.4. **Nilpotent pieces in $\mathfrak{sp}(2n)^*$ and $\mathfrak{o}(2n+1)^*$.** Suppose that $G = Sp(V,\langle,\rangle)$ (resp. $SO(V,Q)$) as in Subsection 2.2. Let $\mathrm{c} \in \underline{\mathcal{N}_{\mathfrak{g}^*}}$ and $\xi \in \mathrm{c}$. Let $V_* = (V_{\geq a})_{a\in\mathbb{Z}}$ be the canonical filtration of V associated to ξ as in [L5, X4], where $V_{\geq a+1} \subset V_{\geq a} \subset V$. If $p \neq 2$, let T_ξ be defined as in Subsection 2.2 (resp. by $\beta(T_\xi v, w) = \beta_\xi(v,w)$ for all $v, w \in V$), then

(a) $V_{\geq a} = \sum_{j\geq \max(0,a)} T^j_\xi(\ker T_\xi^{2j-a+1})$.

The definitions of V_* when $p=2$ are recalled in 2.4.1 (resp. 2.4.3). We define

$$f_a = \dim V_{\geq a}/V_{\geq a+1}.$$

Then $f_a \neq 0$ for finitely many $a \in \mathbb{Z}$ and $f_{-a} = f_a$. The sequence of numbers $(f_a)_{a\in\mathbb{N}}$ ($\mathbb{N} = \{0,1,2,\ldots\}$) depends only on c and not on the choice of $\xi \in \mathrm{c}$; we denote this sequence by Υ_c. Two sequences $(f_a)_{a\in\mathbb{N}}$ and $(h_a)_{a\in\mathbb{N}}$ are equal iff $f_a = h_a$ for all $a \in \mathbb{N}$.

Lemma ([L5, X4])**.** *The orbits* $\mathrm{c}_1, \mathrm{c}_2 \in \underline{\mathcal{N}_{\mathfrak{g}^*}}$ *lie in the same piece if and only if* $\Upsilon_{\mathrm{c}_1} = \Upsilon_{\mathrm{c}_2}$.

Note that if $p \neq 2$, the orbit of $\xi \in \mathcal{N}_{\mathfrak{g}^*}$ is characterized by the partition λ given by the sizes of Jordan blocks of T_ξ. It follows from (a) that

(a′) *if* $p \neq 2$, $\mathrm{c} = \lambda$ *and* $\Upsilon_\mathrm{c} = (f_a)_{a\in\mathbb{N}}$, *then* $f_a = \sum_{i\in\mathbb{N}} m_\lambda(a+2i+1)$ *for all* $a \in \mathbb{N}$.

Moreover each nilpotent piece consists of one orbit when $p \neq 2$.

2.4.1. *Canonical filtrations* V_* *for* $\xi \in \mathcal{N}_{\mathfrak{sp}(2n)^*}$ *(*$p=2$*).* Assume that $G = Sp(V,\langle,\rangle)$ and $p=2$. Let $\xi \in \mathcal{N}_{\mathfrak{g}^*}$ and let $\alpha_\xi, \beta_\xi, T_\xi$ be defined for ξ as in Subsection 2.2. The canonical filtration $V_* = (V_{\geq a})$ associated to ξ is defined by induction on $\dim V$ as follows ([L5]), where $V_{\geq a} = V^\perp_{\geq 1-a}$. If $\xi = 0$, we set $V_{\geq a} = 0$ for all $a \geq 1$ and $V_{\geq a} = V$ for all $a \leq 0$. Hence V_* is defined when $\dim V \leq 1$. Assume now that $\xi \neq 0$ and $\dim V \geq 2$. Let e be the smallest integer such that $T^e_\xi V = 0$, f the smallest integer such that $\alpha_\xi(T^f_\xi V) = 0$ and

$$N = \max(e-1, 2f-1).$$

Then $N \geq 1$. We set

$V_{\geq a} = V$ for all $a \leq -N$; $V_{\geq a} = 0$ for all $a \geq N+1$;

$$V_{\geq -N+1} = \begin{cases} \{v \in V | T^{e-1}_\xi v = 0\} & \text{if } e = 2f+1 \\ \{v \in V | T^{e-1}_\xi v = 0, \alpha_\xi(T^{f-1}_\xi v) = 0\} & \text{if } e = 2f \\ \{v \in V | \alpha_\xi(T^{f-1}_\xi v) = 0\} & \text{if } e < 2f \end{cases};\ V_{\geq N} = V^\perp_{\geq -N+1}.$$

Let $V' = V_{\geq -N+1}/V_{\geq N}$. Then $\langle,\rangle$ induces a nondegenerate symplectic form $\langle,\rangle'$ on V' and α_ξ induces a quadratic form $\alpha_{\xi'}$ corresponding to $\xi' \in \mathcal{N}_{\mathfrak{g}'^*}$, where $\mathfrak{g}' = \mathfrak{sp}(V', \langle,\rangle')$. By induction hypothesis, a canonical filtration $V'_* = (V'_{\geq a})$ of V' is defined for ξ'. For $a \in [-N+1, N]$ we set $V_{\geq a}$ to be the inverse image of $V'_{\geq a}$ under the natural map $V_{\geq -N+1} \to V'$ (note that $V'_{\geq N} = 0$ and $V'_{\geq -N+1} = V'$). This completes the definition of V_*.

2.4.2. Suppose that $p = 2$ and the G-orbit of $\xi \in \mathcal{N}_{\mathfrak{sp}(2n)^*}$ corresponds to $(\lambda, \chi) \in \mathfrak{N}^{*2}_C$. Recall that ([X1]) we have a decomposition $V = \oplus_{a\in[1,r]} W(a)$ of V into mutually orthogonal T_ξ-stable subspaces such that $m_\lambda(i) = \sum_{a\in[1,r]} m_{\lambda^a}(i)$, $\chi(i) = \max_a \chi_a(i)$, where $\alpha_\xi|_{W(a)} = (\lambda^a, \chi_a)$. Moreover, $\alpha_\xi|_{W(a)} = {}^*W_{\chi(\lambda_{2a})}(\lambda_{2a})$, $a \in [1, r]$, where

(a) $\alpha_\xi|_W = {}^*W_l(s)$ means that there exist $v, w \in W$ such that $W = \text{span}\{T^i_\xi v, T^i_\xi w, i \in [0, s-1]\}$, $\langle T^i_\xi v, w\rangle = \delta_{i,s-1}$, $\alpha_\xi(T^i_\xi v) = \delta_{i,l-1}$, $\alpha_\xi(T^i_\xi w) = 0$; we have $\chi_{\alpha_\xi|_W}(i) = \max(0, \min(i - s + l, l))$.

We state some facts which will be used later (see [X1]).

(i) Let W be a T_ξ-stable subspace of V such that $\alpha_\xi|_W = {}^*W_f(e-j)$ with $f > \frac{e-j}{2}$. Let $K_W = \{v \in W | \alpha_\xi(T^{f-1}_\xi v) = 0\}$, $L_W = K^\perp_W \cap W$, $W' = K_W/L_W$ and let $\alpha_{\xi'}$ be the quadratic form on W' induced by α_ξ. Using the basis for W chosen as in (a), one can easily check that

$$\alpha_{\xi'}|_{W'} = {}^*W_{f-1}(e-j-1).$$

(ii) Let W be a T_ξ-stable subspace of V such that $\alpha_\xi|_W = {}^*W_f(e)^a$ (an orthogonal decomposition into a copies of ${}^*W_f(e)$), where $f \leq \frac{e}{2}$. Let $W' = (\ker T^{e-1}_\xi \cap W)/T^{e-1}_\xi W$ and $\alpha_{\xi'}$ be the quadratic form on W' induced by α_ξ. Using the basis for W chosen as in (a), one can easily check that

$$\alpha_{\xi'}|_{W'} = {}^*W_{f-1}(e-2)^a.$$

2.4.3. *Canonical filtrations V_* for $\xi \in \mathcal{N}_{\mathfrak{o}(2n+1)^*}$ ($p = 2$).* Assume that $G = SO(V, Q)$ and $p = 2$. Let $\xi \in \mathcal{N}_{\mathfrak{g}^*}$ and let β_ξ, m, $\{v_i\}$, $\{u_i\}$, W, T_ξ, χ be defined for ξ as in Subsection 2.2. The canonical filtration $V_* = (V_{\geq a})$ associated to ξ is defined by induction on $\dim V$ as follows (see [X4]), where $V_{\geq 1-a} = V^\perp_{\geq a}$ and $Q|_{V_{\geq a}} = 0$ for all $a \geq 1$. If $\xi = 0$ we set $V_{\geq a} = 0$ for all $a \geq 1$ and $V_{\geq a} = V$ for all $a \leq 0$. Hence V_* is defined when $\dim V \leq 1$. Assume now that $\xi \neq 0$ and $\dim V \geq 2$. Let e be the smallest integer such that $T^e_\xi W = 0$, $f = \chi(e)$ and

$$N = \max(2m, m + f - 1).$$

Then $N \geq 1$. We set

$$V_{\geq a} = V \text{ for all } a \leq -N;\ V_{\geq a} = 0 \text{ for all } a \geq N+1;\ V_{\geq N} = V^\perp_{\geq -N+1} \cap Q^{-1}(0),$$

$$V_{\geq -N+1} = \begin{cases} \text{span}\{v_m\} \oplus \ker T^{e-1}_\xi & \text{if } m = 0, \\ \text{span}\{v_i, i \in [0,m], u_i, i \in [1, m-1]\} \oplus W & \text{if } m \geq f, \\ \text{span}\{v_i, i \in [0,m], u_i, i \in [1, m-1]\} & \\ \oplus\{w \in W | Q(T^{f-1}_\xi w) = 0\} & \text{if } e - f < m < f, \\ \text{span}\{v_i, i \in [0,m], u_i, i \in [1, m-1]\} & \\ \oplus\{w \in W | T^{e-1}_\xi w = 0, Q(T^{f-1}_\xi w) = 0\} & \text{if } 0 < m = e - f = f - 1, \\ & \text{or } 0 < m = e - f < f - 1 \text{ and } \rho \neq 0, \\ \text{span}\{v_i, i \in [0,m], u_i, i \in [1, m-1]\} & \\ \oplus\text{span}\{u_0 + w_{**}\} \oplus \ker T^{e-1}_\xi & \text{if } 0 < m = e - f < f - 1 \text{ and } \rho = 0, \end{cases}$$

where $\rho : \ker T_\xi^{e-1} \to \mathbf{k}$ is the map $w \mapsto Q(T_\xi^{f-1}w)$ and $w_{**} \in W$ is such that $\beta(T_\xi^{e-1}w_{**}, w)^2 = Q(T_\xi^{f-1}w)$ for all $w \in W$.

Let $V' = V_{\geq -N+1}/V_{\geq N}$. Then Q induces a non-degenerate quadratic form Q' on V' and β_ξ induces a symplectic form $\beta_{\xi'}$ corresponding to $\xi' \in \mathcal{N}_{\mathfrak{g}'^*}$, where $\mathfrak{g}' = \mathfrak{o}(V', Q')$. By induction hypothesis, a canonical filtration $V'_* = (V'_{\geq a})$ of V' is defined for ξ'. For $a \in [-N+1, N]$ we set $V_{\geq a}$ to be the inverse image of $V'_{\geq a}$ under the natural map $V_{\geq -N+1} \to V'$ (note that $V'_{\geq N} = 0$ and $V'_{\geq -N+1} = V'$). This completes the definition of V_*.

3. Springer fibers

Suppose that $p = 2$ and G is of type B or C in this section.

3.1. For a Borel subgroup B of G, we denote by $\mathfrak{b}$ the Lie algebra of B and define $\mathfrak{n}^* = \{\xi \in \mathfrak{g}^* \mid \xi(\mathfrak{b}) = 0\}$. For $\xi \in \mathcal{N}_{\mathfrak{g}^*}$, denote by $cl_G(\xi)$ the G-orbit of ξ and by $Z_G(\xi)$ the centralizer of ξ in G. Let $\mathcal{B}^G$ be the variety of all Borel subgroups of G and let $\mathcal{B}^G_\xi = \{B \in \mathcal{B}^G \mid \xi \in \mathfrak{n}^*\}$ be the Springer fiber at $\xi \in \mathcal{N}_{\mathfrak{g}^*}$. One can easily adapt the proofs in [St, p. 132] and [S1] to show that $\mathcal{B}^G_\xi$ is connected and all its irreducible components have the same dimension.

Proposition 3.1. *Suppose that $\xi \in \mathcal{N}_{\mathfrak{g}^*}$. We have* $\dim \mathcal{B}^G_\xi = \dfrac{\dim Z_G(\xi) - \textit{rank } G}{2}$.

The proofs of the proposition for $G = Sp(2n)$ and $G = SO(2n+1)$ are given in Subsection 3.2 and Subsection 3.3 respectively.

3.2. Suppose that $G = Sp(V, \langle , \rangle)$ with $\dim V = 2n$ in this subsection. We can identify the variety $\mathcal{B}^G$ with the set $\mathcal{F}^G$ of all complete flags $0 = V_0 \subset V_1 \subset \cdots \subset V_{2n} = V$ such that $V_{2n-i} = V_i^\perp$ and $\langle , \rangle|_{V_i} = 0$ for all $i \leq n$.

Lemma 3.2. *Suppose that $\xi \in \mathcal{N}_{\mathfrak{g}^*}$. We can identify $\mathcal{B}^G_\xi$ with the set*

$$\mathcal{F}^G_\xi = \{F = (V_0 \subset V_1 \subset \cdots \subset V_{2n}) \in \mathcal{F}^G \mid \beta_\xi(V_i, V_{2n+1-i}) = 0 \textit{ and } \alpha_\xi(V_i) = 0 \textit{ for all } i \leq n\}.$$

Proof. Let $\xi \in \mathfrak{sp}(2n)^*$, $x \in \mathfrak{sp}(2n)$ and $F = (V_0 \subset V_1 \subset \cdots \subset V_{2n}) \in \mathcal{F}^G$. There exists a basis e_i, $i \in [-n, n] - \{0\}$, of V such that $\langle e_i, e_j \rangle = \delta_{i+j,0}$ and $V_k = \text{span}\{e_i,\ i \in [1, k]\}$ for all $k \leq n$. Assume that $xe_i = \sum_j x_{ij}e_j$. We have $x_{i,-j} + x_{j,-i} = 0$ and

$$\xi(x) = \sum_{i,j\in[1,n]} x_{ji}\beta_\xi(e_i, e_{-j}) + \sum_{1\leq i<j\leq n} x_{j,-i}\beta_\xi(e_{-j}, e_{-i}) + \sum_{1\leq i<j\leq n} x_{-i,j}\beta_\xi(e_j, e_i)$$
$$+ \sum_{i\in[1,n]} x_{i,-i}\alpha_\xi(e_{-i}) + \sum_{i\in[1,n]} x_{-i,i}\alpha_\xi(e_i).$$

Let $B = \{g \in G | gV_i = V_i\}$ be the Borel subgroup corresponding to F. For $x \in \mathfrak{b}$, we have $x_{ij} = 0$ for $1 \leq i < j \leq n$, and $x_{i,-j} = 0$ for $i, j \in [1, n]$. Hence $\xi \in \mathfrak{n}^*$ if and only if $\beta_\xi(e_i, e_{-j}) = 0$ for all $1 \leq i \leq j \leq n$, $\beta_\xi(e_j, e_i) = 0$ for all $1 \leq i < j \leq n$, and $\alpha_\xi(e_i) = 0$ for all $i \in [1, n]$. The lemma follows. ☐

Let $\xi \in \mathcal{N}_{\mathfrak{g}^*}$ and suppose that $cl_G(\xi) = (\lambda, \chi) \in \mathfrak{N}^{*2}_{C_n}$. Let $F = (V_i) \in \mathcal{F}^G_\xi$. Then we have $\alpha_\xi|_{V_1} = 0$ and $\beta_\xi(V_1, V) = 0$. Let $V' = V_1^\perp/V_1$. The non-degenerate symplectic form $\langle , \rangle$ on V

induces a non-degenerate symplectic form $\langle,\rangle'$ on V'; we write $\mathfrak{g}' = \mathfrak{sp}(V', \langle,\rangle')$. Moreover the quadratic form α_ξ induces a quadratic form $\alpha_{\xi'}$ on V' which corresponds to an orbit (λ', χ') in $\mathcal{N}_{\mathfrak{g}'^*}$. Let $K = \{v \in V | \alpha_\xi(v) = 0, \beta_\xi(v, V) = 0\}$ and for each $(\lambda', \chi') \in \mathfrak{N}^{*2}_{C_{n-1}}$ that arises in this way, let

$$Y_{(\lambda',\chi')} = \{V_1 \in \mathbb{P}(K) | \text{ the quadratic form } \alpha'_\xi \text{ on } V_1^\perp/V_1 \text{ induced by } \alpha_\xi \text{ corresponds to } (\lambda', \chi')\},$$
$$X_{(\lambda',\chi')} = \{F = (V_i) \in \mathcal{F}^G_\xi | V_1 \in Y_{(\lambda',\chi')}\}.$$

Then the fibers of the morphism

$$X_{(\lambda',\chi')} \to Y_{(\lambda',\chi')},\ F = (V_i) \mapsto V_1,$$

are isomorphic to $\mathcal{F}^{G'}_{\xi'}$, where $G' = Sp(2n-2)$ and the G'-orbit of ξ' is (λ', χ'). Now we have that $\dim \mathcal{F}^G_\xi = \max \dim X_{(\lambda',\chi')}$ and by induction hypothesis that $\dim \mathcal{F}^{G'}_{\xi'} = (\dim Z_{G'}(\xi') - n + 1)/2$. For ξ corresponding to (λ, χ), we have $\dim Z_G(\xi) = \sum_{i\geq 1}(i\lambda_i - \chi(\lambda_i))$ (see [X1]). It is then easy to check that $\dim \mathcal{F}^G_\xi = (\dim Z_G(\xi) - n)/2$ using the following lemma ([X3, 5.3]).

Lemma 3.3. *We have* $\dim X_{(\lambda',\chi')} = \dim \mathcal{F}^G_\xi$ *if and only if* (λ', χ') *and* (λ, χ) *are related as follows:* $\lambda'_j = \lambda_j$ *and* $\chi'(\lambda'_j) = \chi(\lambda_j)$ *for* $j \notin \{i-1, i\}$, $\lambda'_{i-1} = \lambda'_i = \lambda_i - 1$, *and* $\chi'(\lambda'_{i-1}) = \chi'(\lambda'_i) \in \{\chi(\lambda_i), \chi(\lambda_i)-1\}$ *satisfies* $[\lambda'_i/2] \leq \chi'(\lambda'_i) \leq \lambda'_i$, $\chi(\lambda_{i+1}) \leq \chi'(\lambda'_i) \leq \chi(\lambda_{i+1}) + \lambda_i - \lambda_{i+1} - 1$ *(this can happen if* $\lambda_{i-1} = \lambda_i > \lambda_{i+1}$*). Moreover* $\dim Y_{(\lambda,\chi')} = i - 1$ *if* $\chi'(\lambda'_i) = \chi(\lambda_i)$ *and* $\dim Y_{(\lambda,\chi')} = i - 2$ *if* $\chi'(\lambda'_i) = \chi(\lambda_i) - 1$.

3.3. Suppose that $G = SO(V, Q)$ with $\dim V = 2n + 1$ in this subsection. We can identify the variety $\mathcal{B}^G$ with the set $\mathcal{F}^G$ of all complete flags $0 = V_0 \subset V_1 \subset \cdots \subset V_{2n+1} = V$ such that $V_{2n+1-i} = V_i^\perp$ and $Q|_{V_i} = 0$ for all $i \leq n$.

Lemma 3.4. *Suppose that* $\xi \in \mathcal{N}_{\mathfrak{g}^*}$*. We can identify* $\mathcal{B}^G_\xi$ *with the set*

$$\mathcal{F}^G_\xi = \{F = (V_0 \subset V_1 \subset \cdots \subset V_{2n+1}) \in \mathcal{F}^G \,|\, \beta_\xi(V_i, V_{2n+2-i}) = 0 \textit{ for all } i \leq n+1\}.$$

Proof. Let $\xi \in \mathfrak{o}(2n+1)^*$, $x \in \mathfrak{o}(2n+1)$ and $F = (V_0 \subset V_1 \subset \cdots \subset V_{2n+1}) \in \mathcal{F}^G$. There exists a basis e_i, $i \in [-n, n]$, of V such that $\beta(e_i, e_j) = \delta_{i+j,0} + \delta_{i,0}\delta_{j,0}$, $Q(e_i) = \delta_{i,0}$, and $V_k = \operatorname{span}\{e_i,\ i \in [1, k]\}$ for all $k \leq n$. Assume that $xe_i = \sum_j x_{ij}e_j$. We have $x_{ij} + x_{-j,-i} = 0$ for all $i, j \in [-n, n] - \{0\}$, $x_{i,-i} = 0$ and $x_{0,i} = 0$ for all $i \in [-n, n]$, and

$$\begin{aligned}\xi(x) = \sum_{i,j\in[1,n]} x_{ji}\beta_\xi(e_i, e_{-j}) + \sum_{1\leq i<j\leq n} x_{j,-i}\beta_\xi(e_{-j}, e_{-i}) + \sum_{1\leq i<j\leq n} x_{-i,j}\beta_\xi(e_j, e_i)\\ + \sum_{i\in[1,n]} x_{i,0}\beta_\xi(e_0, e_{-i}) + \sum_{i\in[1,n]} x_{-i,0}\beta_\xi(e_0, e_i).\end{aligned}$$

Let $B = \{g \in G \,|\, gV_i = V_i\}$ be the Borel subgroup corresponding to F. For $x \in \mathfrak{b}$, we have $x_{ij} = 0$ for $1 \leq i < j \leq n$, $x_{i,-j} = 0$ for $i, j \in [1, n]$, and $x_{i,0} = 0$ for $i \in [1, n]$. Hence $\xi \in \mathfrak{n}^*$ if and only if $\beta_\xi(e_i, e_{-j}) = 0$ for all $1 \leq i \leq j \leq n$, $\beta_\xi(e_j, e_i) = 0$ for all $1 \leq i < j \leq n$, and $\beta_\xi(e_0, e_i) = 0$ for all $i \in [1, n]$. The lemma follows. □

Suppose that $\xi \in \mathcal{N}^*_\mathfrak{g}$ and that $cl_G(\xi) = (m; (\lambda, \chi)) \in \mathfrak{N}^{*2}_{B_n}$. Let $F = (V_i) \in \mathcal{F}^G_\xi$. Then we have $Q|_{V_1} = 0$ and $\beta_\xi(V_1, V) = 0$. Let $V' = V_1^\perp/V_1$. The non-degenerate quadratic form Q on V induces a non-degenerate quadratic form Q' on V'; we write $\mathfrak{g}' = \mathfrak{o}(V', Q')$. The symplectic form β_ξ induces a quadratic form $\beta_{\xi'}$ on V', which corresponds to an orbit $(m'; (\lambda', \chi'))$ in $\mathcal{N}_{\mathfrak{g}'^*}$. Let

$K=\{v\in V|Q(v)=0,\beta_\xi(v,V)=0\}$ and for each $(m';(\lambda',\chi'))\in\mathfrak{N}^{*2}_{B_{n-1}}$ that arises in this way, let

$$\begin{aligned}Y_{m;(\lambda',\chi')}=\{V_1\in\mathbb{P}(K)|&\text{ the symplectic form }\beta'_\xi\text{ on }V_1^\perp/V_1\text{ induced by }\beta_\xi\\&\text{corresponds to }(m';(\lambda',\chi'))\},\\X_{m';(\lambda',\chi')}=\{F=(V_i)&\in\mathcal{F}^G_\xi|V_1\in Y_{m';(\lambda',\chi')}\}.\end{aligned}$$

Then the fibers of the morphism

$$X_{m;(\lambda',\chi')}\to Y_{m';(\lambda',\chi')},\ F=(V_i)\mapsto V_1,$$

are isomorphic to $\mathcal{F}^{G'}_{\xi'}$, where $G'=SO(2n-1)$ and the G'-orbit of ξ' is $(m';(\lambda',\chi'))$. For ξ corresponding to $(m;(\lambda,\chi))$ we have $\dim Z_G(\xi)=m+\sum_{i\geq 1}((i+1)\lambda_i-\chi(\lambda_i))$ (see [X1]). Using the same argument as in the case of $G=Sp(2n)$ and the following lemma ([X3, 6.3]), one can easily check that $\dim\mathcal{F}^G_\xi=(\dim Z_G(\xi)-n)/2$.

Lemma 3.5. *We have* $\dim X_{m',(\lambda',\chi')}=\dim\mathcal{F}^G_\xi$ *if and only if* $(m';(\lambda',\chi'))$ *and* $(m;(\lambda,\chi))$ *are related as follows.*

(a) $m'=m-1$, $\lambda'_i=\lambda_i$ *and* $\chi'(\lambda'_i)=\chi(\lambda_i)$ *(this can happen if* $m-1\geq\lambda_1-\chi(\lambda_1)$*). We have* $\dim Y_{m;(\lambda',\chi')}=0$*;*

(b) $m'=m$, $\lambda'_j=\lambda_j$ *and* $\chi'(\lambda'_j)=\chi(\lambda_j)$ *for* $j\notin\{i-1,i\}$, $\lambda'_{i-1}=\lambda'_i=\lambda_i-1$, *and* $\chi'(\lambda'_{i-1})=\chi'(\lambda'_i)\in\{\chi(\lambda_i),\chi(\lambda_i)-1\}$ *satisfies* $\lambda'_i/2\leq\chi'(\lambda'_i)\leq\lambda'_i$, $\chi(\lambda_{i+1})\leq\chi'(\lambda'_i)\leq\chi(\lambda_{i+1})+\lambda_i-\lambda_{i+1}-1$ *(this can happen if* $\lambda_{i-1}=\lambda_i>\lambda_{i+1}$*). We have* $\dim Y_{m;(\lambda',\chi')}=i$ *if* $\chi'(\lambda'_i)=\chi(\lambda_i)$ *and* $\dim Y_{m;(\lambda',\chi')}=i-1$ *if* $\chi'(\lambda'_i)=\chi(\lambda_i)-1$.

4. INDUCTION FOR NILPOTENT ORBITS

Suppose that $p=2$ and G is of type B or C in this section.

4.1. In this subsection we recall the notion of j-induction in $W^\wedge$ (see [L4] and the references therein). For $\rho\in W^\wedge$, let b_ρ be the smallest integer i such that ρ appears in the ith symmetric power of the reflection representation of W. For a parabolic subgroup W_J of W and $\rho_1\in W_J^\wedge$, there is a unique $\rho\in W^\wedge$ such that $b_\rho=b_{\rho_1}$ and ρ appears in $\mathrm{Ind}_{W_J}^W\rho_1$; we write $\rho=j_{W_J}^W\rho_1$.

Let $\mathfrak{X}=\cup_{n\geq 0}\mathcal{P}_2(n)$ (where we use the notation that $\mathcal{P}_2(0)=\{0\}$). Let $W_0=\{1\}$ and for $n\geq 1$ we denote W_n (resp. S_n) a Weyl group of type B_n or C_n (resp. A_{n-1}). Let sgn denote the sign character of S_n. For each $k\geq 1$, we have ([L4])

$$\text{(a)}\qquad j_{W_n\times S_k}^{W_{n+k}}(\tau\boxtimes\mathrm{sgn})=j_k(\tau)\text{ for }\tau=(\mu)(\nu)\in\mathcal{P}_2(n)\simeq W_n^\wedge,$$

where $j_k:\mathfrak{X}\to\mathfrak{X},\ (\mu)(\nu)\mapsto(\mu')(\nu')$, is defined by

$$\mu'_i=\begin{cases}\mu_i+1&\text{if }i\leq(k+1)/2\\\mu_i&\text{otherwise}\end{cases},\ \nu'_i=\begin{cases}\nu_i+1&\text{if }i\leq k/2\\\nu_i&\text{otherwise.}\end{cases}$$

It is easy to see that j_k is injective and $j_k\circ j_l=j_l\circ j_k$.

4.2. For a parabolic subgroup P of G with a Levi subgroup L and unipotent radical U_P, let $\mathfrak{p}$, $\mathfrak{n}_\mathfrak{p}$, $\mathfrak{l}$ denote the Lie algebra of P, U_P, L respectively. We define $\mathfrak{p}^*=\{\xi\in\mathfrak{g}^*\,|\,\xi(\mathfrak{n}_\mathfrak{p})=0\}$, $\mathfrak{n}_\mathfrak{p}^*=\{\xi\in\mathfrak{g}^*\,|\,\xi(\mathfrak{l}\oplus\mathfrak{n}_\mathfrak{p})=0\}$ and $\mathfrak{l}^*=\{\xi\in\mathfrak{g}^*\,|\,\xi(\mathfrak{n}_\mathfrak{p}\oplus\mathfrak{n}_\mathfrak{p}^-)=0\}$ where $\mathfrak{g}=\mathfrak{l}\oplus\mathfrak{n}_\mathfrak{p}\oplus\mathfrak{n}_\mathfrak{p}^-$.

Let c' be a nilpotent L-orbit in $\mathfrak{l}^*$. Since $\mathfrak{g}^*$ has finitely many nilpotent G-orbits, there exists a unique nilpotent G-orbit c in $\mathfrak{g}^*$ such that $\mathrm{c}\cap(\mathrm{c}'+\mathfrak{n}_{\mathfrak{p}}^*)$ is dense in $\mathrm{c}'+\mathfrak{n}_{\mathfrak{p}}^*$ (note that $\mathrm{c}'+\mathfrak{n}_{\mathfrak{p}}^*\subset\mathcal{N}_{\mathfrak{g}^*}$). Following [LS] we say that c is obtained by inducing c' from $\mathfrak{l}^*$ to $\mathfrak{g}^*$ and denote $\mathrm{c}=\mathrm{Ind}_{\mathfrak{l}^*,\mathfrak{p}^*}^{\mathfrak{g}^*}\mathrm{c}'$.

Proposition 4.1. *Suppose that* $\mathrm{c}'\in\underline{\mathcal{N}_{\mathfrak{l}^*}}$ *and* $\mathrm{c}=\mathit{Ind}_{\mathfrak{l}^*,\mathfrak{p}^*}^{\mathfrak{g}^*}\mathrm{c}'$. *We have* $\gamma_{\mathfrak{g}^*}(\mathrm{c})=j_{W_L}^W(\gamma_{\mathfrak{l}^*}(\mathrm{c}'))$, *where* W_L *is the Weyl group of* L.

The proposition is an analog of [LS, 3.5]. To prove it one can adapt the proof in [S3, 4.1] (another proof in unipotent case is given in [LS]). We outline the proof here. Suppose that $\gamma_{\mathfrak{g}^*}(\mathrm{c})=\rho\in W^\wedge$ and $\gamma_{\mathfrak{l}^*}(\mathrm{c}')=\rho'\in W_L^\wedge$. Let $\xi'\in\mathrm{c}'$ and $\xi\in(\xi'+\mathfrak{n}_{\mathfrak{p}}^*)\cap\mathrm{c}$. One can show that $b_\rho=\dim\mathcal{B}_\xi^G$ and $b_{\rho'}=\dim\mathcal{B}_{\xi'}^L$ by direct computation (using Proposition 3.1 and the known information on $\dim Z_G(\xi)$ and on b_ρ) or by adapting the proof in [S3]. It is easy to adapt the proof in [LS] to show that $\dim Z_G(\xi)=\dim Z_L(\xi')$ and $\dim\mathcal{B}_\xi^G=\dim\mathcal{B}_{\xi'}^L$. It then follows that $b_\rho=b_{\rho'}$. Now let $Y_{\xi,\xi'}=\{g\in G|g^{-1}.\xi\in\xi'+\mathfrak{n}_{\mathfrak{p}}^*\}$ and $I_{\xi,\xi'}$ the set of irreducible components of $Y_{\xi,\xi'}$ of dimension $\frac{1}{2}(\dim Z_G(\xi)+\dim Z_{G'}(\xi'))+\dim U_P$. Then $A_G(\xi)\times A_L(\xi')$ acts on $I_{\xi,\xi'}$; we denote the corresponding representation by $\varepsilon_{\xi,\xi'}$. We have $I_{\xi,\xi'}\neq\emptyset$ and $\langle\rho',\mathrm{Res}_{W_L}^W(\rho)\rangle_{W_L}=\langle 1,\varepsilon_{\xi,\xi'}\rangle_{A_G(\xi)\times A_L(\xi')}\neq 0$ (see [L2, X3]). It now follows from the definition of j-induction that $\rho'=j_{W_L}^W(\rho')$.

Note that it follows from Proposition 4.1 and the injectivity of $\gamma_{\mathfrak{g}^*}$ that $\mathrm{c}=\mathrm{Ind}_{\mathfrak{l}^*,\mathfrak{p}^*}^{\mathfrak{g}^*}\mathrm{c}'$ does not depend on the choice of $P\supset L$. Hence we can write $\mathrm{c}=\mathrm{Ind}_{\mathfrak{l}^*}^{\mathfrak{g}^*}\mathrm{c}'$.

4.3. Let $\tilde{\mathfrak{g}}=\mathfrak{sp}(2n+2k)$ (resp. $\mathfrak{o}(2n+2k+1)$) be the Lie algebra of $\tilde{G}=Sp(2n+2k)$ (resp. $SO(2n+2k+1)$). Let P be a parabolic subgroup of $\tilde{G}$ such that $\mathfrak{l}\cong\mathfrak{sp}(2n)\oplus\mathfrak{gl}(k)$ (resp. $\mathfrak{o}(2n+1)\oplus\mathfrak{gl}(k)$) for a Levi subgroup L of P. Let c be the nilpotent orbit in $\mathfrak{l}^*$ corresponding to the nilpotent orbit $(\lambda,\chi)\in\mathfrak{N}_{C_n}^{*2}$ (resp. $(m;(\lambda,\chi))\in\mathfrak{N}_{B_n}^{*2}$) and to the 0 orbit in $\mathfrak{gl}(k)^*$. Let $(\tilde{\lambda},\tilde{\chi})$ (resp. $(\tilde{m};(\tilde{\lambda},\tilde{\chi}))$) correspond to the orbit $\tilde{\mathrm{c}}=\mathrm{Ind}_{\mathfrak{l}^*}^{\tilde{\mathfrak{g}}^*}\mathrm{c}$. We write $(\tilde{\lambda},\tilde{\chi})=j_k(\lambda,\chi)$ (resp. $(\tilde{m};(\tilde{\lambda},\tilde{\chi}))=j_k(m;(\lambda,\chi))$) in view of the following commutative diagrams (see 4.1 (a) and Proposition 4.1)

$$\begin{tikzcd}
\mathfrak{N}^{*2} \arrow[r, "j_k"] \arrow[d, "\gamma^*"] & \mathfrak{N}^{*2} \arrow[d, "\gamma^*"] \\
\mathfrak{X} \arrow[r, "j_k"] & \mathfrak{X}
\end{tikzcd}$$

where γ^* is the Springer correspondence map. Using this one easily sees that (see also [S2])

(a) *For every* $(\lambda,\chi)\in\mathfrak{N}_C^{*2}$ *(resp.* $(m;(\lambda,\chi))\in\mathfrak{N}_B^{*2}$*), there exists a sequence of integers* $l_1,\ldots,l_s$ *such that* $j_{l_1}\circ\cdots\circ j_{l_s}((\lambda,\chi))$ *(resp.* $j_{l_1}\circ\cdots\circ j_{l_s}(m;(\lambda,\chi))$*) is of the form* $j_{k_1}\circ\cdots\circ j_{k_r}(0)$ *for some sequence* $k_1,\ldots,k_r$.

5. CLOSURE RELATIONS ON NILPOTENT ORBITS IN $\mathfrak{g}^*$ WHEN G IS CLASSICAL

5.1. Assume that G is of type B or C. By identifying $W^\wedge$ with the set of $\mathcal{P}_2(n)$ we get order relations on $W^\wedge$ (see Subsection 2.1). We have

Theorem 5.1. *Suppose that* $p=2$ *and* $\mathrm{c},\mathrm{c}'\in\underline{\mathcal{N}_{\mathfrak{g}^*}}$. *We have* $\mathrm{c}<\mathrm{c}'$ *if and only if* $\gamma_{\mathfrak{g}^*}(\mathrm{c})<\gamma_{\mathfrak{g}^*}(\mathrm{c}')$.

Note that theorem is true when G is of type D by [S2] or when $p\neq 2$ (see [X2]) (in these cases we can identify $\mathfrak{g}^*$ with $\mathfrak{g}$). We prove the theorem in the remainder of this section (where we assume that $p=2$) using similar arguments as in [S2] most of the time. In the reduction process of 5.3 we use a different argument without using the theory of sheets and packets.

5.2. Assume that $G=Sp(V,\langle,\rangle)$ (resp. $SO(V,Q)$). Let V_1 and V_2 be orthogonal subspaces of V such that $V=V_1\oplus V_2$. Then the restrictions of $\langle,\rangle$ (resp. Q) on V_1 and V_2 are non-degenerate. Let $G_1=Sp(V_1,\langle,\rangle|_{V_1})$ (resp. $G_1=SO(V_1,Q|_{V_1})$) and $G_2=Sp(V_2,\langle,\rangle|_{V_2})$ (resp. $G_2=SO(V_2,Q|_{V_2})$). Using the isomorphism $\mathfrak{g}^*\xrightarrow{\sim}\mathfrak{Q}(V)$ (resp. $\mathfrak{g}^*\xrightarrow{\sim}\mathfrak{S}(V)$) (see 2.2 (a)) we have a natural inclusion $\mathfrak{g}_1^*\oplus\mathfrak{g}_2^*\subset\mathfrak{g}^*$.

Lemma 5.2. *If $\xi_1,\eta_1\in\mathcal{N}_{\mathfrak{g}_1^*}$ and $\xi_2,\eta_2\in\mathcal{N}_{\mathfrak{g}_2^*}$ are such that $cl_{G_1}(\xi_1)\leq cl_{G_1}(\eta_1)$ and $cl_{G_2}(\xi_2)\leq cl_{G_2}(\eta_2)$, then $cl_G(\xi_1+\xi_2)\leq cl_G(\eta_1+\eta_2)$.*

5.3. For $\tau\in\mathfrak{X}_R^{*2}$, we denote $\mathrm{c}_\tau\in\mathfrak{N}_R^{*2}$ the corresponding orbit, where R stands for B or C (type of G). We show by induction on $\dim G$ that if $\tau'\leq\tau$ then $\mathrm{c}_{\tau'}\leq\mathrm{c}_\tau$ (see Subsection 5.4 for $G=Sp(2n)$ and Subsection 5.5 for $G=SO(2n+1)$). We may assume that this is true for classical groups of strictly smaller dimension and that $\tau'<\tau$, $\{\tau''\in\mathfrak{X}_R^{*2}|\tau'<\tau''<\tau\}=\emptyset$. We have the following reduction process.

Reduction 1. Suppose in the case of $G=SO(2n+1)$ (resp. $Sp(2n)$), we have $\mathfrak{g}^*\supset\mathfrak{g}_1^*\oplus\mathfrak{g}_2^*$ with $0\neq\mathfrak{g}_1^*\neq\mathfrak{g}^*$ and $\mathfrak{g}_1^*\cong\mathfrak{o}(2k+1)^*$, $\mathfrak{g}_2^*\cong\mathfrak{o}(2n-2k)^*$ (resp. $\mathfrak{g}_1^*\cong\mathfrak{sp}(2k)^*$, $\mathfrak{g}_2^*\cong\mathfrak{sp}(2n-2k)^*$). Assume that we can find $\xi=\xi_1+\xi_2\in\mathrm{c}_\tau$, $\xi'=\xi_1'+\xi_2'\in\mathrm{c}_{\tau'}$ with $\xi_1,\xi_1'\in\mathfrak{g}_1^*$ corresponding to τ_1,τ_1' in $\mathfrak{X}$ and $\xi_2,\xi_2'\in\mathfrak{g}_2^*$ corresponding to τ_2,τ_2' in $\mathfrak{X}$, such that $\tau_1'\leq\tau_1$ and $\tau_2'\leq\tau_2$. Then by induction hypothesis and Lemma 5.2 we have $\mathrm{c}_{\tau'}\leq\mathrm{c}_\tau$.

Reduction 2. Assume that we can find $\omega>\omega'$ in $\mathfrak{X}_R^{*2}$ such that $\tau=j_{i_1}\circ\cdots\circ j_{i_r}(\omega)$ and $\tau'=j_{i_1}\circ\cdots\circ j_{i_r}(\omega')$ for some positive integers $i_1,\ldots,i_r$. Then by induction hypothesis and the fact that induction for orbits preserves order, we have $\mathrm{c}_{\tau'}\leq\mathrm{c}_\tau$.

5.4. Assume that $G=Sp(2n)$. We have $\mathfrak{X}_C^{*2}=\{(\mu)(\nu)|\nu_i\leq\mu_i+1\}$. Let $\tau=(\mu)(\nu),\tau'=(\mu')(\nu')\in\mathfrak{X}_{C_n}^{*2}$ be such that $\tau>\tau'$ and $\{\tau''\in\mathfrak{X}_{C_n}^{*2}|\tau>\tau''>\tau'\}=\emptyset$.

Lemma 5.3. *Suppose that reduction 1 does not apply. One of the following is true:*

(a) $\tau=(\mu_1)(\nu_1)$, $\tau'=(\mu_1-1)(\nu_1+1)$;

(b) $\tau=(\mu_1,\mu_2)(\nu_1,\nu_2)$, $\tau'=(\mu_1,\mu_2+1)(\nu_1-1,\nu_2)$.

Proof. Note that we can apply Reduction 1 if $\mu_1^*\geq 2$ or $\nu_1^*\geq 2$, and for some j, $\mu_j+\nu_j=\mu_j'+\nu_j'$ with $\mu_j\geq\mu_j'$ or $\nu_j'=\nu_k'$ for some $k<j$. We denote by (1^a) the partition with all parts 1 and multiplicity a. Since Reduction 1 does not apply, if $\mu_1+\nu_1=\mu_1'+\nu_1'$, then we have $\mu_1^*\leq 1,\nu_1^*\leq 1$ and thus τ,τ' are as in case (a). From now on we assume that $\mu_1+\nu_1>\mu_1'+\nu_1'$. Let $r=\mu_1$ and $a=\mu_r^*$. Then $r>0$ (otherwise $\tau=(0)(1^n)$ is minimal). We have the following cases.

1) $\mu_1>\mu_1'$. Let $b=\nu_r^*$ and $c=\nu_{r+1}^*$. Then $c\leq a$ and $b\leq\mu_{r-1}^*$.

i) If $b<a$, then there exists $\tau''\in\mathfrak{X}_C^{*2}$ such that $\tau=\tau''+(1^a)(1^b)$. One easily verifies that $\tau>\tau''+(1^b)(1^a)\geq\tau'$ (note that $\mu_i-1=\mu_1-1\geq\mu_i'$ for $i\in[b+1,a]$). Hence $\tau'=\tau''+(1^b)(1^a)$.

ii) If $b\geq a$ and $c\neq 0$, then there exists $\tau''\in\mathfrak{X}_C^{*2}$ such that $\tau=\tau''+(1^a)(1^c)$. One easily verifies that $\tau>\tau''+(1^{a+1})(1^{c-1})\geq\tau'$ (note that $\nu_c-1=\mu_1\geq\nu_c'$ and $\nu_i+\mu_{i+1}-1=2\mu_1-1\geq\nu_i'+\mu_{i+1}'$ for $i\in[c+1,a-1]$). Hence $\tau'=\tau''+(1^{a+1})(1^{c-1})$.

iii) If $b\geq a$, $c=0$ and $b<\mu_{r-1}^*$, then there exists $\tau''\in\mathfrak{X}_C^{*2}$ such that $\tau=\tau''+(1^a)(1^b)$. One easily verifies that $\tau > \tau'' + (1^{b+1})(1^{a-1}) \geq \tau'$ (note that $\mu_a + \nu_a - 1 = 2\mu_1 - 1 \geq \mu'_a + \nu'_a$ and $\nu_i + \mu_i = 2\mu_1 - 1 \geq \nu'_i + \mu'_i$ for $i \in [a+1, b]$). Hence $\tau' = \tau'' + (1^{b+1})(1^{a-1})$.

iv) If $b \geq a$, $c = 0$ and $b = \mu^*_{r-1}$, then let $\tau'' = (\mu'')(\nu'')$ with $\nu'' = \nu$ and $\mu''_i = \mu_i$ except that $\mu''_a = \mu_a - 1$ and $\mu''_{b+1} = \mu_{b+1} + 1$. Then one easily verifies that $\tau > \tau'' \geq \tau'$ (note that $\mu_a - 1 = \mu_1 - 1 \geq \mu'_a$, $\mu_a + \nu_a - 1 = 2\mu_1 - 1 \geq \mu'_a + \nu'_a$ and $\mu_i \geq \mu'_i, \nu_i \geq \nu'_i$ for $i \in [a+1, b]$). Hence $\tau' = \tau''$.

Since $\mu_1 > \mu'_1$, in case (i) we have $b = 0$, cases (ii) and (iii) do not happen, in case (iv) we have $a = 1$. As Reduction 1 does not apply, in case (i) we have $a = 1$, $\tau = (\mu_1)(\nu_1)$ and $\tau' = (\mu_1 - 1)(\nu_1 + 1)$; in case (iv) we have $b = 1$, $\tau = (\mu_1, \mu_2)(\mu_1, \nu_2)$ and $\tau' = (\mu_1 - 1, \mu_2 + 1)(\mu_1, \nu_2)$, but notice that $\tau > (\mu_1, \mu_2 + 1)(\mu_1 - 1, \nu_2) > \tau'$ ($\nu_2 \leq \mu_1 - 1$ since $b = 1$) so case (iv) does not happen. It follows that τ and τ' are as in (a).

2) $\mu_1 = \mu'_1$. Then $\nu_1 > \nu'_1$. Let $s = \nu_1$ and $b = \nu^*_s$.

i) If $a \leq b$, then there exists $\tau'' \in \mathfrak{X}^{*2}_C$ such that $\tau = \tau'' + (1^a)(1^b)$. One easily verifies that $\tau > \tau'' + (1^{b+1})(1^{a-1}) \geq \tau'$ (note that $\nu_i - 1 = \nu_1 - 1 \geq \nu'_i$ for $i \in [a, b]$). Hence $\tau' = \tau'' + (1^{b+1})(1^{a-1})$.

ii) If $a > b$ and $\nu_a = s - 1$, then there exists $\tau'' \in \mathfrak{X}^{*2}_C$ such that $\tau = \tau'' + (1^a)(1^b)$. One easily verifies that $\tau > \tau'' + (1^{a+1})(1^{b-1}) \geq \tau'$ (note that $\nu_b - 1 = \nu_1 - 1 \geq \nu'_b$, $\nu_i = \nu_1 - 1 \geq \nu'_i$ and $\mu_i = \mu_1 \geq \mu'_i$ for $i \in [b+1, a]$). Hence $\tau' = \tau'' + (1^{a+1})(1^{b-1})$.

iii) If $a > b$, $\nu_a \leq s - 2$, then there exists $b' \in [b+1, a-1]$ such that $\nu_{b+1} = \cdots = \nu_{b'} = s - 1 > \nu_{b'+1}$. Let $\tau'' = (\mu'')(\nu'')$ with $\mu'' = \mu$ and $\nu''_i = \nu_i$ except that $\nu''_b = \nu_b - 1$ and $\nu''_{b'+1} = \nu_{b'+1} + 1$. Then one easily verifies that $\tau > \tau'' \geq \tau'$ (note that $\nu_b - 1 = \nu_1 - 1 \geq \nu'_b$, and $\mu_i = \mu_1 \geq \mu'_i, \nu_i = \nu_1 - 1 \geq \nu'_i$ for $i \in [b+1, b']$). Hence $\tau' = \tau''$.

Since $\nu_1 > \nu'_1$, we have that in case (i) $a = 1$, in cases (ii) and (iii) $b = 1$. As Reduction 1 does not apply, in case (i) we have $b = 1$, $\tau = (\mu_1, \mu_2)(\nu_1, \nu_2)$ and $\tau' = (\mu_1, \mu_2 + 1)(\nu_1 - 1, \nu_2)$; case (ii) does not happen ($a \geq 2$); in case (iii) we have $\nu_2 \leq \nu_1 - 2$, $\tau = (\mu_1, \mu_1)(\nu_1, \nu_2)$ and $\tau' = (\mu_1, \mu_1)(\nu_1 - 1, \nu_2 + 1)$ (but notice that $\tau > (\mu_1, \mu_1 - 1)(\nu_1, \nu_2 + 1) > \tau'$, so this case does not happen). It follows that τ and τ' are as in (b). □

Proposition 5.4. *Suppose that Reductions 1 and 2 do not apply. One of the following is true:*

(i) $\tau = (1)(0)$, $\tau' = (0)(1)$;
(ii) $\tau = (1)(1)$, $\tau' = (1,1)(0)$;
(iii) $\tau = (1)(2)$, $\tau' = (1,1)(1)$;
(iv) $\tau = (1)(2,1)$, $\tau' = (1,1)(1,1)$;
(v) $\tau = (\frac{n}{2}, \frac{n}{2} - 1)(1)$, $\tau' = (\frac{n}{2}, \frac{n}{2})(0)$.

Proof. In case (a) of Lemma 5.3 we have $\tau = j_1^{\mu_1 - \nu_1 - 1} \circ j_2^{\nu_1}(\omega)$ and $\tau' = j_1^{\mu_1 - \nu_1 - 1} \circ j_2^{\nu_1}(\omega')$, where $\omega = (1)(0) > \omega' = (0)(1)$. In case (b) of Lemma 5.3 we have $\tau = j_1^{\mu_1 - \mu_2 - 1} \circ j_3^{\nu_1 - \nu_2 - 1} \circ j_4^{\nu_2}(\omega)$ and $\tau' = j_1^{\mu_1 - \mu_2 - 1} \circ j_3^{\nu_1 - \nu_2 - 1} \circ j_4^{\nu_2}(\omega')$ if $\nu_1 \leq \mu_2 + 1$, where $\omega = (\mu_2 - \nu_1 + 2, \mu_2 - \nu_1 + 1)(1) > \omega' = (\mu_2 - \nu_1 + 2, \mu_2 - \nu_1 + 2)(0)$; $\tau = j_1^{\mu_1 - \nu_1} \circ j_2^{\nu_1 - \mu_2 - 1} \circ j_3^{\mu_2 - \nu_2} \circ j_4^{\nu_2}(\omega)$ and $\tau' = j_1^{\mu_1 - \nu_1} \circ j_2^{\nu_1 - \mu_2 - 1} \circ j_3^{\mu_2 - \nu_2} \circ j_4^{\nu_2}(\omega')$ if $\mu_2 + 2 \leq \nu_1 \leq \mu_1$ and $\nu_2 \leq \mu_2$, where $\omega = (1)(1) > \omega' = (1,1)(0)$; $\tau = j_2^{\nu_1 - \mu_2 - 2} \circ j_3^{\mu_2 - \nu_2} \circ j_4^{\nu_2}(\omega)$ and $\tau' = j_2^{\nu_1 - \mu_2 - 2} \circ j_3^{\mu_2 - \nu_2} \circ j_4^{\nu_2}(\omega')$ if $\nu_1 = \mu_1 + 1$ and $\nu_2 \leq \mu_2$, where $\omega = (1)(2) > \omega' = (1,1)(1)$; $\tau = j_1^{\mu_1 - \nu_1 + 1} \circ j_2^{\nu_1 - \mu_2 - 2} \circ j_4^{\mu_2}(\omega)$ and $\tau' = j_1^{\mu_1 - \nu_1 + 1} \circ j_2^{\nu_1 - \mu_2 - 2} \circ j_4^{\mu_2}(\omega')$ if $\nu_2 = \mu_2 + 1$ and $\nu_1 \geq \mu_2 + 2$, where $\omega = (1)(2,1) > \omega' = (1,1)(1,1)$. Since Reduction 2 does not apply, the proposition follows. □

It remains to show that in each case of Proposition 5.4 we have $\mathrm{c}_{\tau'} < \mathrm{c}_\tau$. In case (i) it is obvious that $\mathrm{c}_{\tau'} < \mathrm{c}_\tau$ as τ' corresponds to the 0 orbit. In each case (ii)-(v), we choose an element $\xi \in \mathrm{c}_\tau$ and define a family $\{g_t \in Sp(V),\ t \in \mathbf{k}^*\}$ such that $\lim_{t\to 0} g_t^{-1}.\xi = \xi' \in \mathrm{c}_{\tau'}$. Then it follows that $\mathrm{c}_{\tau'} < \mathrm{c}_\tau$. The elements ξ (or equivalently the quadratic form α_ξ associated to ξ) and the family $\{g_t \in Sp(V),\ t \in \mathbf{k}^*\}$ in each case are given as follows. We have $\alpha_{g.\xi}(v) = \alpha_\xi(g^{-1}v)$.

Let $e_i, i \in [-n,n] - \{0\}$ be a basis of V such that $\langle e_i, e_j\rangle = \delta_{i+j,0}$.

(ii) $\alpha_\xi(\sum a_ie_i) = a_{-2}^2 + a_1a_{-2}$; $g_te_1 = te_1,\ g_te_2 = e_2,\ g_te_{-1} = \frac{1}{t}e_{-1},\ g_te_{-2} = e_{-2}$. Then $\alpha_{g_t^{-1}.\xi}(\sum a_ie_i) = a_{-2}^2 + ta_1a_{-2}$.

(iii) $\alpha_\xi(\sum a_ie_i) = a_1^2 + a_1a_{-2} + a_2a_{-3}$; $g_te_1 = e_1,\ g_te_2 = \frac{1}{t}e_2,\ g_te_3 = \frac{1}{t}e_3, g_te_{-1} = e_{-1},\ g_te_{-2} = te_{-2},\ g_te_{-3} = te_{-3}$. Then $\alpha_{g_t^{-1}.\xi}(\sum a_ie_i) = a_1^2 + a_2a_{-3} + ta_1a_{-2}$.

(iv) $\alpha_\xi(\sum a_ie_i) = a_1a_{-2} + a_2a_{-3}$; $g_te_1 = te_1,\ g_te_2 = e_2 + e_4,\ g_te_3 = e_3,\ g_te_4 = te_4,\ g_te_{-1} = \frac{1}{t}e_{-1},\ g_te_{-2} = e_{-2},\ g_te_{-3} = e_{-3},\ g_te_{-4} = \frac{1}{t}(e_{-4} + e_{-2})$. Then $\alpha_{g_t^{-1}.\xi}(\sum a_ie_i) = a_1a_{-4} + a_2a_{-3} + ta_1a_{-2}$.

(v) $\alpha_\xi(\sum a_ie_i) = \sum_{i\in[1,d]} a_ia_{-i-1} + \sum_{i\in[d+2,2d-1]} a_ia_{-i-1} + a_d^2 + a_{2d}^2$, where $d = \frac{n}{2}$; $g_te_1 = e_1,\ g_te_{-1} = e_{-1},\ g_te_i = \frac{1}{t}(e_i + e_{i+d})$ and $g_te_{-i} = te_{-i}$ for $i \in [2,d],\ g_te_{d+1} = \frac{1}{t}e_{d+1},\ g_te_{-d-1} = te_{-d-1},\ g_te_i = e_i$ and $g_te_{-i} = e_{-i} + e_{-i+d}$ for $i \in [d+2, 2d]$. Then $\alpha_{g_t^{-1}.\xi}(\sum a_ie_i) = a_1a_{-d-2} + \sum_{i\in[2,d]} a_ia_{-i-1} + \sum_{i\in[d+2,2d-1]} a_ia_{-i-1} + a_{2d}^2 + ta_1a_{-2}$.

5.5. Assume that $G = SO(2n+1)$. We have $\mathfrak{X}_B^{*2} = \{(\mu)(\nu) | \nu_i \geq \mu_{i+1}\}$. Let $\tau = (\mu)(\nu), \tau' = (\mu')(\nu') \in \mathfrak{X}_{B_n}^{*2}$ be such that $\tau > \tau'$ and $\{\tau'' \in \mathfrak{X}_{B_n}^{*2} | \tau > \tau'' > \tau'\} = \emptyset$.

Proposition 5.5. *Suppose that reduction 1 and 2 do not apply. Then* $\tau = (1)(n-1)$, $\tau' = (0)(n)$.

Proof. Note that we can apply Reduction 1 if $\mu_1 = \mu_1'$ or if for some $j \geq 1$, $\mu_{j+1} + \nu_j = \mu'_{j+1} + \nu'_j$ with $\nu_j \geq \nu'_j$ or $\mu'_{j+1} = \mu'_k$ for some $k < j+1$.

Since Reduction 1 does not apply, we have $\mu_1 > \mu_1'$. Let $r = \mu_1$ and $a = \mu_1^*$. Let $\tau'' = (\mu'')(\nu'')$ with $\mu_i'' = \mu_i - 1$ for $i \in [1,a]$ and $\nu_i'' = \nu_i + 1$ for $i \in [1,a]$. Then one easily verifies that $\tau'' \in \mathfrak{X}_B^{*2}$ and $\tau > \tau'' \geq \tau'$. Hence $\tau' = \tau''$. Since Reduction 1 does not apply, we have $a = 1$, $\tau = (\mu_1, \mu_2)(\nu_1)$ and $\tau' = (\mu_1 - 1, \mu_2)(\nu_1 + 1)$. We have $\tau = j_1^{\mu_1-\nu_1-1} \circ j_2^{\nu_1-\mu_2} \circ j_3^{\mu_2}((1)(0))$ and $\tau' = j_1^{\mu_1-\nu_1-1} \circ j_2^{\nu_1-\mu_2} \circ j_3^{\mu_2}((0)(1))$ if $\mu_1 \geq \nu_1 + 2$; $\tau = j_2^{\mu_1-\mu_2-1} \circ j_3^{\mu_2}((1)(\nu_1 + 1 - \mu_1))$ and $\tau' = j_2^{\mu_1-\mu_2-1} \circ j_3^{\mu_2}((0)(\nu_1 + 2 - \mu_1))$ if $\mu_1 \leq \nu_1 + 1$. Since Reduction 2 does not apply, τ and τ' are as in the proposition. ☐

Let $\tau = (1)(n-1)$ and $\tau = (0)(n)$. We show that $\mathrm{c}_\tau > \mathrm{c}_{\tau'}$. If $n = 1$, this is obvious. Assume that $n \geq 2$. Let e_i, $i \in [-n,n]$ be a basis of V such that $Q(e_i) = \delta_{i,0}$ and $\langle e_i, e_j\rangle = \delta_{i+j,0} - \delta_{i,0}\delta_{j,0}$. Let β_ξ be the symplectic form corresponding to $\xi \in \mathcal{N}_{\mathfrak{g}^*}$ such that $\beta_\xi(\sum_{i\in[-n,n]} a_ie_i, \sum_{i\in[-n,n]} b_ie_i) = a_0b_{-1} + b_0a_{-1} + a_nb_{-1} + b_na_{-1} + \sum_{i\in[2,n-1]}(a_ib_{-i-1} + b_ia_{-i-1})$. Then $\xi \in \mathrm{c}_\tau$. Let $g_t \in SO(V)$, $t \in \mathbf{k}^*$ be defined by:

$$g_te_0 = e_0,\ g_te_i = \frac{1}{t}e_i,\ g_te_{-i} = te_{-i},\ i \in [1,n].$$

We have $\beta_{g_t^{-1}.\xi}(\sum a_ie_i, \sum b_ie_i) = \beta_\xi(g_t(\sum a_ie_i), g_t(\sum b_ie_i)) = t(a_0b_{-1} + b_0a_{-1}) + a_nb_{-1} + b_na_{-1} + \sum_{i\in[2,n-1]}(a_ib_{-i-1} + b_ia_{-i-1})$. Thus $\lim_{t\to 0} g_t^{-1}.\xi \in \mathrm{c}_{\tau'}$ and $\mathrm{c}_{\tau'} < \mathrm{c}_\tau$.

5.6. We show that if $\mathrm{c}_{\tau'} \leq \mathrm{c}_\tau$ then $\tau' \leq \tau$. Since induction for orbits preserves order and $j_k(\tau') \leq j_k(\tau)$ iff $\tau' \leq \tau$, we may use the operation j_k as often as needed. We show that

$$(*) \qquad \sum_{j\in[1,i]} (\mu'_j + \nu'_j) \leq \sum_{j\in[1,i]} (\mu_j + \nu_j),$$

$$(**) \qquad \sum_{j\in[1,i-1]} (\mu'_j + \nu'_j) + \mu'_i \leq \sum_{j\in[1,i-1]} (\mu_j + \nu_j) + \mu_i.$$

We write $\mathrm{codim}(\tau,\tau') = \mathrm{codim}_{\overline{\mathrm{c}_\tau}}\mathrm{c}_{\tau'}$ for $\tau,\tau' \in \mathfrak{X}_R^{*2}$. For each $l \geq 1$, let $\Gamma'_l = \{(\tau,\tau') \in \mathfrak{X}_R^{*2} \times \mathfrak{X}_R^{*2},\ \mathrm{c}_\tau \geq \mathrm{c}_{\tau'}$ and $(*)$ fails for $i = l\}$, $\Delta'_l = \{(\tau,\tau') \in \mathfrak{X}_R^{*2} \times \mathfrak{X}_R^{*2},\ \mathrm{c}_\tau \geq \mathrm{c}_{\tau'}$ and $(**)$ fails for $i = l\}$, $d = \min\{\mathrm{codim}(\tau,\tau')|(\tau,\tau') \in \Gamma'_l\}$, $d' = \min\{\mathrm{codim}(\tau,\tau')|(\tau,\tau') \in \Delta'_l\}$ and

$$\Gamma_l = \{(\tau,\tau') \in \Gamma'_l|\mathrm{codim}(\tau,\tau') = d\},\ \Delta_l = \{(\tau,\tau') \in \Delta'_l|\mathrm{codim}(\tau,\tau') = d'\}.$$

It is enough to show that $\Gamma_l = \emptyset$ and $\Delta_l = \emptyset$.

It follows from the definitions that we have

Lemma 5.6. *(a) if $(\tau,\tau') \in \Gamma_l$ (resp. Δ_l) then $(j_k(\tau), j_k(\tau')) \in \Gamma_l$ (resp. Δ_l).*

(b) if $\tau,\tau',\tau'' \in \mathfrak{X}_R^{*2}$, $\mathrm{c}_\tau \geq \mathrm{c}_{\tau''} \geq \mathrm{c}_{\tau'}$ and $(\tau,\tau') \in \Gamma_l$ (or Δ_l), then $\tau = \tau''$ or $\tau'' = \tau'$.

Assume that $G = Sp(V,\langle,\rangle)$ (resp. $SO(V,Q)$). Let $(\tau,\tau') \in \Gamma_l$ or Δ_l. We can assume that $\tau = j_k(\omega)$ for some $\omega \in \mathfrak{X}_R^{*2}$. Let Σ be an isotropic subspace of V with $\dim\Sigma = k$ and let $W \subset V$ be such that $\Sigma^\perp = \Sigma \oplus W$. Let P be the parabolic subgroup that stabilizes Σ and L the Levi subgroup of P that stabilizes W. Then $L \cong Sp(2n-2k) \times GL(k)$ (resp. $SO(2n-2k+1) \times GL(k)$). We have $\overline{\mathrm{c}_\tau} = G.(\overline{\mathrm{c}_\omega} + \mathfrak{n}_\mathfrak{p}^*)$. Since $\mathrm{c}_\tau \geq \mathrm{c}_{\tau'}$, there exists $\xi' \in \mathrm{c}_{\tau'}$ such that $\alpha_{\xi'}|_\Sigma = 0$, $\beta_{\xi'}(\Sigma,\Sigma^\perp) = 0$ (resp. $\beta_{\xi'}(\Sigma,\Sigma^\perp) = 0$) and such that $\alpha_{\xi'}$ (resp. $\beta_{\xi'}$) induces a quadratic form (resp. symplectic form) on $\Sigma^\perp/\Sigma \cong W$ which corresponds to an element $\eta' \in \overline{\mathrm{c}_\omega}$. Let $\omega' \in \mathfrak{X}_R^{*2}$ correspond to the $Sp(2n-2k)$-orbit (resp. $SO(2n-2k+1)$-orbit) of η'. We have $\mathrm{c}_{\tau'} \leq \mathrm{Ind}_{\mathfrak{l}^*}^{\mathfrak{g}^*}\mathrm{c}_{\omega'} \leq \mathrm{c}_\tau = \mathrm{Ind}_{\mathfrak{l}^*}^{\mathfrak{g}^*}\mathrm{c}_\omega$ and thus

either a) $\mathrm{c}_{\tau'} = \mathrm{Ind}_{\mathfrak{l}^*}^{\mathfrak{g}^*}\mathrm{c}_{\omega'}$
or b) $\mathrm{c}_{\omega'} = \mathrm{c}_\omega$.

In case a) we have $\tau' = j_k(\omega')$ and $(\omega,\omega') \in \Gamma_l$ or Δ_l. Applying j_k when needed we can assume that $\tau = j_{k_r} \circ \cdots \circ j_{k_1}(0)$ for some sequence $k_1, \ldots, k_r$ (see 4.3 (a)) and moreover:

(i) $k_1 = 2s$ is even;

(ii) $0 \leq k_i - k_{i+1} \leq 1$ for $i \leq r-1$ and $k_i = k_{i+1}$ if $3 \nmid i$;

(iii) ${\mu'_1}^* \leq s,\ {\nu'_1}^* \leq s$.

We apply the previous construction with $k = k_r$, and repeat the argument in case b) with (τ,τ') replaced by (ω,ω') (conditions (i)-(iii) above are still satisfied) until we reach a point that $\omega = \omega'$. Hence we are reduced to the case that $(\tau,\tau') \in \Gamma_l$ (or Δ_l), $\tau = j_{k_r} \circ \cdots \circ j_{k_1}(0)$, $\omega = \omega' = j_{k_{r-1}} \circ \cdots \circ j_{k_1}(0)$ (τ,τ' satisfy (i)-(iii) above), where τ' and ω' are related as in the above construction. We will study more closely the relation between τ' and ω' and derive a contradiction.

Suppose that $\omega = \omega' = (\sigma)(\delta)$. Let $a = [(k_r+1)/2]$, $b = [k_r/2]$. We show that (see 5.7 and 5.8)

c) $\mu'_i \leq \sigma_1 + 1 = \mu_a$ for all i;

d) $\mu'_i + \nu'_i \leq \sigma_1 + \delta_1 + 2 = \mu_b + \nu_b$ for all i;

e) if $a < s$, then $\mu'_i \geq \sigma_i = \mu_i$ and $\nu'_i \geq \delta_i = \nu_i$ for $i \in [a+1, s]$ (resp. if $b < s$, then $\mu'_{i+1} \geq \sigma_{i+1} = \mu_{i+1}$ and $\nu'_i \geq \delta_i = \nu_i$ for $i \in [b+1, s]$).

If $(\tau, \tau') \in \Gamma_l$, it follows from c) and d) that $l \geq b+1$ and from e) that $l \leq b$; if $(\tau, \tau') \in \Delta_l$ it follows from c) and d) that $l \geq a+1$ and from e) that $l \leq a$. This is the required contradiction.

Suppose that $r > 1$. We have if $k_{r-1} = k_r$ is odd, then $a = b+1$, $\sigma = (r-1)^a(3i_2)(3i_4)\cdots$, $\delta = (r-1)^b(3i_1)(3i_3)\cdots$; if $k_{r-1} = k_r$ is even, then $a = b$, $\sigma = (r-1)^a(3i_1)(3i_3)\cdots$, $\delta = (r-1)^b(3i_2)(3i_4)\cdots$; if $k_{r-1} = k_r + 1$ is even, then $3|r-1$, $a = b+1$, $\sigma = (r-1)^a(3i_1)(3i_3)\cdots$, $\delta = (r-1)^{b+1}(3i_2)(3i_4)\cdots$; if $k_{r-1} = k_r + 1$ is odd, then $3|r-1$, $a = b$, $\sigma = (r-1)^{a+1}(3i_2)(3i_4)\cdots$, $\delta = (r-1)^b(3i_1)(3i_3)\cdots$; where in each case $r-1 > 3i_1$ and $i_1 > i_2 > i_3 > \cdots$.

5.7. We prove 5.6 c)-e) for $G = Sp(V, \langle , \rangle)$ by studying the quadratic form $\alpha_{\eta'}$ on $\Sigma^\perp/\Sigma$ induced by $\alpha_{\xi'}$. Let $c = \max(\mu_1'^*, \nu_1'^*)$.

If $c = 1$, then V is indecomposable (as $T_{\xi'}$-module), and $\alpha_{\xi'} = {}^*W_{\chi(\lambda_1)}(\lambda_1)$ (where $\chi(\lambda_1) = \mu_1'$, $\lambda_1 = \mu_1' + \nu_1'$; see 2.4.2 (a)). Then one easily verifies that $(\Sigma^\perp/\Sigma, \alpha_{\eta'}) \cong {}^*W_{\chi(\lambda_1)-1}(\lambda_1 - 2)$ (if $\dim\Sigma = 2$), ${}^*W_{\chi(\lambda_1)-1}(\lambda_1 - 1)$ or ${}^*W_{\chi(\lambda_1)}(\lambda_1 - 1)$ (if $\dim\Sigma = 1$). Thus ${\sigma_1}^* \leq 1, {\delta_1}^* \leq 1$ and $\mu_1' - 1 \leq \sigma_1 \leq \mu_1'$, $\nu_1' - 1 \leq \delta_1 \leq \nu_1'$; in this case (c) and (d) hold.

If $c > 1$, then V is decomposable, and there exists an orthogonal decomposition $V = \oplus_{i=1}^c W_i$ with W_i indecomposable, $\dim W_1 \geq \dim W_2 \geq \cdots \geq \dim W_c$, such that $\Sigma = \oplus_{i=1}^c (W_i \cap \Sigma)$. We apply the previous result for each factor in the decomposition $\Sigma^\perp/\Sigma = \oplus_{i=1}^c (W_i \cap \Sigma^\perp)/(W_i \cap \Sigma)$. Assume that $\alpha_{\xi'}|_{W_i} = {}^*W_{l_i'}(\lambda_i')$ (where $\lambda_i' = \mu_i' + \nu_i'$). Then $l_i' \leq \mu_i'$ and $l_j' = \mu_1'$ for some j. Suppose that $\alpha_{\eta'}|_{(W_i \cap \Sigma^\perp)/(W_i \cap \Sigma)} = {}^*W_{l_i''}(\lambda_i'')$. We have $\sigma_1 = \max\{l_i''\} \geq \max\{l_i' - 1\} = \mu_1' - 1$ and $\sigma_1 + \delta_1 = \max\{\lambda_i''\} \geq \max\{\lambda_i' - 2\} = \mu_1' + \nu_1' - 2$; thus c) and d) hold.

Now we prove e). Assume that $a < s$. Then $r > 1$ and $c = s$. Note that $\sigma_i + \delta_i - (\sigma_{i+1} + \delta_{i+1}) \geq 3$ for $i \in [a, s-1]$; hence if ψ is a permutation such that $\dim(W_{\psi(i)} \cap \Sigma^\perp)/(W_{\psi(i)} \cap \Sigma) \geq \dim(W_{\psi(i+1)} \cap \Sigma^\perp)/(W_{\psi(i+1)} \cap \Sigma)$, then $\psi(i) = i$ for $i \in [a+1, s]$. Moreover since $\sigma_{i+1} - \sigma_{i+2} \geq 3$ and $\delta_i - \delta_{i+1} \geq 3$ for $i \in [a, s-1]$, $l_i'' = \sigma_i$ for all $i \in [a+1, s]$. It follows that $l_i'' = r - 1 = \sigma_i$ for $i \in [1, a]$. Now we have $l_i' \geq l_i'' = \sigma_i > \sigma_{i+1} + 1 = l_{i+1}'' + 1 \geq l_{i+1}'$ and $\lambda_i' - l_i' \geq \delta_i > \delta_{i+1} + 1 \geq \lambda_{i+1}' - l_{i+1}'$ for $i \in [a+1, s-1]$, $\lambda_a' - l_a' \geq \lambda_a'' - l_a'' = \delta_j \geq \delta_a > \delta_{a+1} + 1 \geq \lambda_{a+1}' - l_{a+1}'$ for some $j \in [1, a]$; hence $l_i' = \mu_i'$ for $i \in [a+1, s]$. Then e) follows.

5.8. We prove 5.6 c)-e) for $G = SO(V, Q)$ by studying the symplectic form $\beta_{\eta'}$ on $\Sigma^\perp/\Sigma$ induced by $\beta_{\xi'}$. Let $c = \max(\mu_1'^* - 1, \nu_1'^*)$.

If $c = 0$, then $\mu_1' = m$, $\nu_1' = 0$, $V = \text{span}\{v_i, i \in [0, m], u_i, i \in [0, m-1]\}$ (where v_i, u_i are as in 2.2 (b) (c)), $\Sigma = \text{span}\{v_0\}$ and thus $\sigma_1^* \leq 1$, $\sigma_1 = m - 1, \delta_1 = 0$; in this case (c) and (d) hold.

If $c > 0$, then there exists an orthogonal decomposition $V = \text{span}\{v_i, i \in [0, m], u_i, i \in [0, m-1]\} \oplus_{i=1}^c W_i$ with W_i indecomposable (as $T_{\xi'}$-module), $\dim W_1 \geq \dim W_2 \geq \cdots \geq \dim W_s$, such that $\Sigma = (\text{span}\{v_i, i \in [0, m], u_i, i \in [0, m-1]\} \cap \Sigma) \oplus_{i=1}^c (W_i \cap \Sigma)$. Assume that $T_{\xi'}|_{W_i} = W_{l_i'}(\lambda_i')$ (where $\lambda_i' = \mu_{i+1}' + \nu_i'$; notation as in [X2, 5.6]). One easily verifies that $T_{\eta'}|_{(W_i \cap \Sigma^\perp)/(W_i \cap \Sigma)} = W_{l_i'-1}(\lambda_i' - 2)$ (if $\dim(W_i \cap \Sigma) = 2$), $W_{l_i'-1}(\lambda_i' - 1)$ or $W_{l_i'}(\lambda_i' - 1)$ (if $\dim(W_i \cap \Sigma) = 1$). Write $T_{\eta'}|_{(W_i \cap \Sigma^\perp)/(W_i \cap \Sigma)} = W_{l_i''}(\lambda_i'')$. Then $l_i' - 1 \leq l_i'' \leq l_i'$ and $\lambda_i'' \geq \lambda_i' - 2$. We have $\mu_1' - 1 \leq \sigma_1 \leq \mu_1'$, and $\delta_1 = \max\{l_i'', \lambda_1'' - \sigma_1\} \geq \max\{l_i' - 1, \mu_2' + \nu_1' - 2 - \sigma_1\}$; thus $\delta_1 \geq \nu_1' - 2$ if $\sigma_1 = \mu_1'$ and $\delta_1 \geq \nu_1' - 1$ if $\sigma_1 = \mu_1' - 1$ (note that $\nu_1' = \max\{l_i', \mu_2' + \nu_1' - \mu_1'\}$). Hence c) and d) follow.

We prove e). Assume that $b < s$. Then $r > 1$ and $c = s$. Note that $\sigma_{i+1} + \delta_i - (\sigma_{i+2} + \delta_{i+1}) \geq 3$ for $i \in [b, s-1]$; hence if ψ is a permutation such that $\dim(W_{\psi(i)} \cap \Sigma^\perp)/(W_{\psi(i)} \cap \Sigma) \geq \dim(W_{\psi(i+1)} \cap \Sigma^\perp)/(W_{\psi(i+1)} \cap \Sigma)$, then $\psi(i) = i$ for $i \in [b+1, s]$. Moreover for all $i \in [b+1, s]$, $l_i'' = \delta_i$ since

$\sigma_i - \sigma_{i+1} \geq 3$ and $\delta_i - \delta_{i+1} \geq 3$. It follows that $l''_i = r - 1 = \delta_i$ for all $i \in [1, b]$. Now we have $l'_i \geq l''_i = \delta_i > \delta_{i+1} + 1 = l''_{i+1} + 1 \geq l'_{i+1}$ and $\lambda'_i - l'_i \geq \sigma_{i+1} > \sigma_{i+2} + 1 \geq \lambda'_{i+1} - l'_{i+1}$ for $i \in [b+1, s-1]$, $\lambda'_b - l'_b \geq \lambda''_b - l''_b = \sigma_{j+1} \geq \sigma_{b+1} > \sigma_{b+2} + 1 \geq \lambda'_{b+1} - l'_{b+1}$ for some $j \in [1, b]$ or (if $b = 0$) $m \geq \sigma_{b+1} > \sigma_{b+2} + 1 \geq \lambda'_{b+1} - l'_{b+1}$; hence $l'_i = \nu'_i \geq \delta_i$ and $\mu'_{i+1} = \lambda'_i - l'_i \geq \sigma_{i+1}$ for $i \in [b+1, s]$. This completes the proof of Theorem 5.1.

6. NILPOTENT PIECES IN $\mathfrak{g}^*$ FOR TYPE B, C

Assume that $p = 2$ and G is of type B or C throughout this section unless otherwise stated.

6.1. Suppose that $G = Sp(2n)$ (resp. $SO(2n+1)$). Let $\mathfrak{X}^1_R \subset \mathfrak{X}$ denote the image of the Springer correspondence map $\gamma_{G_\mathbb{C}} : \underline{\mathcal{U}_{G_\mathbb{C}}} \to W^\wedge$, where as before R stands for B or C (the type of G or $G_\mathbb{C}$). We have (see [L1, X2])

$$\mathfrak{X}^1_C = \{(\mu)(\nu) | \mu_{i+1} - 1 \leq \nu_i \leq \mu_i + 1\},\ \mathfrak{X}^1_B = \{(\mu)(\nu) \in | \mu_{i+1} \leq \nu_i \leq \mu_i + 2\}.$$

Let $\tilde{\tau} \in \mathfrak{X}^1_R$ and let $\mathrm{c} \in \underline{\mathcal{N}_{\mathfrak{g}^*}}$ be such that $\gamma_{\mathfrak{g}^*}(\mathrm{c}) = \tilde{\tau}$. Define $\Sigma^{\mathfrak{g}^*}_{\tilde{\tau}}$ to be the set of all orbits $\mathrm{c}' \in \underline{\mathcal{N}_{\mathfrak{g}^*}}$ such that $\mathrm{c}' \leq \mathrm{c}$ and $\mathrm{c}' \nleq \mathrm{c}''$ for any $\mathrm{c}'' < \mathrm{c}$ with $\gamma_{\mathfrak{g}^*}(\mathrm{c}'') \in \mathfrak{X}^1_R$. We show that

(a) $\{\Sigma^{\mathfrak{g}^*}_{\tilde{\tau}}\}_{\tilde{\tau} \in \mathfrak{X}^1_R}$ *form a partition of* $\mathcal{N}_{\mathfrak{g}^*}$.

Following [X2], we define maps

$$\Phi_R : \mathfrak{X}^{*2}_R \to \mathfrak{X}^1_R,\ (\mu)(\nu) \mapsto (\tilde{\mu})(\tilde{\nu}),$$

where if $R = B$, then

$$\tilde{\mu}_i = \begin{cases} [\frac{\mu_i + \nu_i - 1}{2}] & \text{if } \nu_i > \mu_i + 2 \\ \mu_i & \text{if } \nu_i \leq \mu_i + 2 \end{cases},\quad \tilde{\nu}_i = \begin{cases} [\frac{\mu_i + \nu_i + 2}{2}] & \text{if } \nu_i > \mu_i + 2 \\ \nu_i & \text{if } \nu_i \leq \mu_i + 2 \end{cases},\ i \geq 1;$$

if $R = C$, then $\tilde{\mu}_1 = \mu_1$ and

$$\tilde{\mu}_{i+1} = \begin{cases} [\frac{\mu_{i+1} + \nu_i + 1}{2}] & \text{if } \nu_i < \mu_{i+1} - 1 \\ \mu_{i+1} & \text{if } \nu_i \geq \mu_{i+1} - 1 \end{cases},\ \tilde{\nu}_i = \begin{cases} [\frac{\mu_{i+1} + \nu_i}{2}] & \text{if } \nu_i < \mu_{i+1} - 1 \\ \nu_i & \text{if } \nu_i \geq \mu_{i+1} - 1 \end{cases},\ i \geq 1.$$

It is easy to verify that in each case we get a well-defined element $(\tilde{\mu})(\tilde{\nu}) \in \mathfrak{X}^1_R$. We have

(b) *if* $(\tilde{\mu}')(\tilde{\nu}') \in \mathfrak{X}^1_R$, $(\mu)(\nu) \in \mathfrak{X}^{*2}_R$, *then* $\Phi_R((\mu')(\nu')) = (\mu')(\nu')$, $(\mu)(\nu) \leq \Phi_R((\mu)(\nu))$; *if moreover* $(\mu)(\nu) \leq (\tilde{\mu}')(\tilde{\nu}')$, *then* $\Phi_R((\mu)(\nu)) \leq (\tilde{\mu}')(\tilde{\nu}')$.

In fact, if $R = C$, (b) follows from [X2, 4.2]; if $R = B$, one can prove (b) by the same argument. Now in view of Theorem 5.1, it follows from the definition of $\Sigma^{\mathfrak{g}^*}_{\tilde{\tau}}$ and (b), (c) that for each $\tilde{\tau} \in \mathfrak{X}^1_R$,

(d) $\gamma_{\mathfrak{g}^*}(\Sigma^{\mathfrak{g}^*}_{\tilde{\tau}}) = \Phi^{-1}_R(\tilde{\tau})$.

Then (a) follows from (d).

6.2. We define a map

$$\Psi^*_R : \mathfrak{N}^{*2}_R \to \underline{\mathcal{U}}_{G_\mathbb{C}},\ \mathrm{c} \mapsto \tilde{\lambda} = (\tilde{\lambda}_1 \geq \tilde{\lambda}_2 \geq \cdots)$$

as follows such that each fiber is an nilpotent piece (see Proposition 6.1).

Assume that $G = Sp(2n)$ and $\mathrm{c} = (\lambda, \chi) \in \mathfrak{N}^{*2}_C$. If $\chi(\lambda_1) > \frac{\lambda_1}{2}$, then $\tilde{\lambda}_1 = 2\chi(\lambda_1)$, if $\chi(\lambda_{2i}) > \frac{\lambda_{2i}}{2}$ and $\chi(\lambda_{2i}) > \chi(\lambda_{2i+1})$, then

$$\tilde{\lambda}_{2i} = \begin{cases} \lambda_{2i} - \chi(\lambda_{2i}) + \chi(\lambda_{2i+1}) & \text{if } \chi(\lambda_{2i}) - \lambda_{2i} + \chi(\lambda_{2i+1}) \geq 1 \\ 2(\lambda_{2i} - \chi(\lambda_{2i})) & \text{if } \chi(\lambda_{2i}) - \lambda_{2i} + \chi(\lambda_{2i+1}) \leq 0 \end{cases},$$

if $\chi(\lambda_{2i+1}) > \frac{\lambda_{2i+1}}{2}$ and $\lambda_{2i+1} - \chi(\lambda_{2i+1}) < \lambda_{2i} - \chi(\lambda_{2i})$, then

$$\tilde{\lambda}_{2i+1} = \begin{cases} \lambda_{2i} - \chi(\lambda_{2i}) + \chi(\lambda_{2i+1}) & \text{if } \chi(\lambda_{2i}) - \lambda_{2i} + \chi(\lambda_{2i+1}) \geq 1 \\ 2\chi(\lambda_{2i+1}) & \text{if } \chi(\lambda_{2i}) - \lambda_{2i} + \chi(\lambda_{2i+1}) \leq 0 \end{cases};$$

otherwise $\tilde{\lambda}_i = \lambda_i$.

Assume that $G = SO(2n+1)$ and $\mathrm{c} = (m; (\lambda, \chi)) \in \mathfrak{N}_B^{*2}$. Let

$$\tilde{\lambda}_1 = \begin{cases} m + \chi(\lambda_1) & \text{if } \chi(\lambda_1) \geq m+2 \\ 2m+1 & \text{if } \chi(\lambda_1) < m+2 \end{cases}, \quad \tilde{\lambda}_2 = \begin{cases} m + \chi(\lambda_1) & \text{if } \chi(\lambda_1) \geq m+2 \\ 2\chi(\lambda_1) - 1 & \text{if } [\frac{\lambda+1}{2}] < \chi(\lambda_1) < m+2 \\ \lambda_1 & \text{if } \chi(\lambda_1) \leq [\frac{\lambda+1}{2}] \end{cases}.$$

For $i \geq 1$, if $\chi(\lambda_{2i}) > \frac{\lambda_{2i}}{2}$ and $\chi(\lambda_{2i}) > \chi(\lambda_{2i+1})$, then

$$\tilde{\lambda}_{2i+1} = \begin{cases} \lambda_{2i} - \chi(\lambda_{2i}) + \chi(\lambda_{2i+1}) & \text{if } \chi(\lambda_{2i}) - \lambda_{2i} + \chi(\lambda_{2i+1}) \geq 2 \\ 2(\lambda_{2i} - \chi(\lambda_{2i})) + 1 & \text{if } \chi(\lambda_{2i}) - \lambda_{2i} + \chi(\lambda_{2i+1}) \leq 1 \end{cases},$$

if $\chi(\lambda_{2i+1}) > \frac{\lambda_{2i+1}}{2}$ and $\lambda_{2i+1} - \chi(\lambda_{2i+1}) < \lambda_{2i} - \chi(\lambda_{2i})$, then

$$\tilde{\lambda}_{2i+2} = \begin{cases} \lambda_{2i} - \chi(\lambda_{2i}) + \chi(\lambda_{2i+1}) & \text{if } \chi(\lambda_{2i}) - \lambda_{2i} + \chi(\lambda_{2i+1}) \geq 2 \\ 2\chi(\lambda_{2i+1}) - 1 & \text{if } \chi(\lambda_{2i}) - \lambda_{2i} + \chi(\lambda_{2i+1}) \leq 1 \end{cases};$$

otherwise $\tilde{\lambda}_{2i+1} = \lambda_{2i}$, $\tilde{\lambda}_{2i+2} = \lambda_{2i+1}$.

We show that

(a) $\gamma_{G_{\mathbb{C}}} \circ \Psi_R^* = \Phi_R \circ \gamma_{\mathfrak{g}^*}$.

Let $G = SO(2n+1)$ (resp. $Sp(2n)$) and $\mathrm{c} \in \mathfrak{N}_R^{*2}$. Assume that $\gamma_{\mathfrak{g}^*}(\mathrm{c}) = (\mu)(\nu)$, $\Phi_R((\mu)(\nu)) = (\tilde{\mu})(\tilde{\nu})$ and $(\tilde{\mu})(\tilde{\nu}) = \gamma_{G_{\mathbb{C}}}(\tilde{\lambda})$. Using the definition of Φ_R, one easily shows that $\tilde{\nu}_i = \tilde{\mu}_i + 2$ iff $\nu_i \geq \mu_i + 2$ and $\mu_i + \nu_i$ is even, and $\tilde{\nu}_i = \tilde{\mu}_{i+1}$ iff $\nu_i = \mu_{i+1}$ (resp. $\tilde{\nu}_i = \tilde{\mu}_i + 1$ iff $\nu_i = \mu_i + 1$, and $\tilde{\nu}_i = \tilde{\mu}_{i+1} - 1$ iff $\nu_i \leq \mu_{i+1} - 1$ and $\nu_i + \mu_{i+1}$ is odd). Using this and the description of the map $\gamma_{G_{\mathbb{C}}}$ (see [X2, 2.4]), one easily verifies that

$$\tilde{\lambda}_{2i-1} = \begin{cases} \mu_i + \nu_i & \text{if } \mu_i \leq \nu_i - 2 \\ 2\mu_i + 1 & \text{if } \nu_i - 2 < \mu_i < \nu_{i-1} \\ 2\mu_i & \text{if } \mu_i = \nu_{i-1} \end{cases}, \quad \tilde{\lambda}_{2i} = \begin{cases} \mu_i + \nu_i & \text{if } \nu_i \geq \mu_i + 2 \\ 2\nu_i - 1 & \text{if } \mu_{i+1} < \nu_i < \mu_i + 2 \\ 2\nu_i & \text{if } \nu_i = \mu_{i+1} \end{cases}$$

$$\left(\text{resp. } \tilde{\lambda}_{2i-1} = \begin{cases} \mu_i + \nu_{i-1} & \text{if } \mu_i \geq \nu_{i-1} + 1 \\ 2\mu_i & \text{if } \nu_i \leq \mu_i \leq \nu_{i-1} \\ 2\mu_i + 1 & \text{if } \mu_i = \nu_i - 1 \end{cases}, \quad \tilde{\lambda}_{2i} = \begin{cases} \mu_{i+1} + \nu_i & \text{if } \nu_i \leq \mu_{i+1} - 1 \\ 2\nu_i & \text{if } \mu_{i+1} \leq \nu_i \leq \mu_i \\ 2\nu_i - 1 & \text{if } \nu_i = \mu_i + 1 \end{cases} \right).$$

It is then easy to verify that $\tilde{\lambda} = \Psi_R^*(\mathrm{c})$ using the description of $\gamma_{\mathfrak{g}^*}$ (see 2.3).

Proposition 6.1. *Two orbits* $\mathrm{c}_1, \mathrm{c}_2 \in \underline{\mathcal{N}_{\mathfrak{g}^*}}$ *lie in the same nilpotent piece as defined in* [L5, X4] *if and only if* $\Psi_R^*(\mathrm{c}_1) = \Psi_R^*(\mathrm{c}_2)$.

Note that the proposition computes the nilpotent pieces in $\mathfrak{g}^*$ explicitly. Now in view of (a) and 6.1 (d), it follows from Proposition 6.1 that each set $\Sigma_{\tilde{\tau}}^{\mathfrak{g}^*}$ is a nilpotent piece defined in [L5, X4]. One can also define a partition of $\mathcal{N}_{\mathfrak{g}^*}$ into special pieces as in [L3, X2] and show that each special piece is a union of nilpotent pieces. The proof of the proposition is given in the remainder of this section following the argument used in [X2].

6.3. Suppose that $G = Sp(V, \langle , \rangle)$ (resp. $SO(V, Q)$). Let $\mathrm{c} \in \underline{\mathcal{N}_{\mathfrak{g}^*}}$ and $\Psi^*_R(\mathrm{c}) = \tilde{\mathrm{c}} = \tilde{\lambda}$. Suppose that $\Upsilon_{\mathrm{c}} = (f_a)_{a\in\mathbb{N}}$ and $\Upsilon_{\tilde{\mathrm{c}}} = (\tilde{f}_a)_{a\in\mathbb{N}}$ (see 2.4.3). We show that

(a) $f_a = \tilde{f}_a$ *for all* $a \in \mathbb{N}$.

Then Proposition 6.1 follows from (a) and Lemma 2.4.3.

We prove (a) by induction on $\dim V$. Let $\xi \in \mathrm{c}$. If $\xi = 0$, (a) is obvious. Assume from now on that $\xi \neq 0$. Let $V_* = (V_{\geq a})$, V', ξ' be associated to ξ and N, e, f defined for ξ as in 2.4.1 (resp. 2.4.3). Let c' be the orbit of ξ' in $\mathfrak{g}'^*$ and let $\tilde{\mathrm{c}}' = \Psi^2_{\mathfrak{g}'^*}(\mathrm{c}') = \tilde{\lambda}'$. Suppose that $\Upsilon_{\mathrm{c}'} = (f'_a)$ and $\Upsilon_{\tilde{\mathrm{c}}'} = (\tilde{f}'_a)$. Since $\dim V' < \dim V$, by induction hypothesis $f'_a = \tilde{f}'_a$ for all $a \in \mathbb{N}$. By the definition of V_* we have that for all $a \in [0, N-1]$, $f_a = f'_a$ and thus $f_a = \tilde{f}'_a$. We show that

$$\text{(b)}\quad \begin{array}{l} \tilde{\lambda}_1 = N+1, m_{\tilde{\lambda}}(\tilde{\lambda}_1) = f_N, m_{\tilde{\lambda}'}(\tilde{\lambda}_1) = 0,\ m_{\tilde{\lambda}'}(\tilde{\lambda}_1 - 2) = m_{\tilde{\lambda}}(\tilde{\lambda}_1 - 2) + m_{\tilde{\lambda}}(\tilde{\lambda}_1), \\ m_{\tilde{\lambda}'}(i) = m_{\tilde{\lambda}}(i) \textit{ for all } i \neq \tilde{\lambda}_1, \tilde{\lambda}_1 - 2. \end{array}$$

It then follows from (b) and 2.4 (a′) that $\tilde{f}_a = 0$ for all $a \geq N+1$, $\tilde{f}_N = f_N$, and that $\tilde{f}_a = \tilde{f}'_a$ for all $a \in [0, N-1]$. Hence (a) follows (note that $f_a = 0$ for all $a \geq N+1$).

The proof of (b) is given in subsections 6.4-6.6 (resp. 6.7-6.9).

6.4. Assume that $G = Sp(V, \langle , \rangle)$ throughout subsection 6.6. We keep the notations in 6.3. Suppose that $\mathrm{c} = (\lambda, \chi)$. We show that

$$\text{(a) } f_N = \begin{cases} 1 & \textit{if } e < 2f, \\ m_\lambda(e) & \textit{if } e = 2f+1, \textit{ or } e = 2f \textit{ and } \chi(e-1) = f-1, \\ m_\lambda(e)+1 & \textit{if } e = 2f \textit{ and } \chi(e-1) = f. \end{cases}$$

Recall that $f_N = \dim V_{\geq N}$ and $V_{\geq N} = V^{\perp}_{\geq -N+1}$ (see 2.4.1). We describe $V_{\geq N}$ in various cases in the following and then (a) follows.

Suppose that $e < 2f$. Then the map $\rho : V \to \mathbf{k},\ v \mapsto \sqrt{\alpha_\xi(T^{f-1}_\xi v)}$ is linear and thus $V_{\geq N} = (\ker \rho)^\perp$ is a line.

Suppose that $e = 2f+1$, or $e = 2f$ and $\chi(e-1) = f-1$. Then $V_{\geq N} = (\ker T^{e-1}_\xi)^\perp = \mathrm{Im} T^{e-1}_\xi$.

Suppose that $e = 2f$ and $\chi(e-1) = f$. Let E be subspace of V such that $V = \ker T^{e-1}_\xi \oplus E$ and let $W = \sum_{i\in[0,e]} T^i_\xi E$. Then $\langle , \rangle|_W$ is non-degenerate (in fact, if $\langle \sum_i T^i_\xi v_i, W\rangle = 0$, where $v_i \in E$, then $\langle T^{e-1}_\xi v_0, V\rangle = 0$ and thus $v_0 \in E \cap \ker T^{e-1}_\xi = 0$; now use induction and similar argument one shows that $v_i = 0$). Thus $V = W \oplus W^\perp$, $W^\perp$ is T_ξ-stable and $T^{e-1}_\xi W^\perp = 0$ ($W \supset T^{e-1}_\xi V$ implies that $W^\perp \subset (\mathrm{Im} T^{e-1}_\xi)^\perp = \ker T^{e-1}_\xi$). We have $V_{\geq -N+1} = (\ker T^{e-1}_\xi \cap W) \oplus \{v \in W^\perp | \alpha_\xi(T^{f-1}_\xi v) = 0\}$ (note that $\ker T^{e-1}_\xi \cap W \subset \mathrm{Im} T_\xi$). Thus $V_{\geq N} = \mathrm{Im} T^{e-1}_\xi \oplus L$, where $L \subset W^\perp$ is a line (we apply the result in the first case for $W^\perp$).

6.5. We describe $\mathrm{c}' = (\lambda', \chi')$ in various cases as follows. Let $j \geq 0$ be the unique integer such that $\chi(e-j) = f$ and $\chi(e-j-1) < f$.

(i) $e = 2f+1$, or $e = 2f$ and $\chi(e-1) = f-1$. We have

$m_{\lambda'}(e) = 0$, $m_{\lambda'}(e-2) = m_\lambda(e) + m_\lambda(e-2)$ *(if* $e > 2$*)*, $m_{\lambda'}(i) = m_\lambda(i)$ *for* $i \notin \{e, e-2\}$, $\chi'(\lambda_i) = \chi(\lambda_i)$ *for* $\lambda_i \notin \{e, e-2\}$, $\chi'(e-2) = f-1$ *if* $\chi(e-2) \leq f-1$ *and* $\chi'(e-2) = f$ *if* $\chi(e-2) = f$.

(ii) $e=2f$ and $\chi(e-1)=f$. We have

$m_{\lambda'}(e)=0,\ m_{\lambda'}(e-2)=m_\lambda(e-2)+m_\lambda(e)+2\delta_{j,1}-2\delta_{j,2}$ (if $e>2$), $m_{\lambda'}(e-j)=m_\lambda(e-j)-2+\delta_{j,2}m_\lambda(e)$, $m_{\lambda'}(e-j-1)=m_\lambda(e-j-1)+2+\delta_{j,1}m_\lambda(e)$ (if $e>j+1$), $m_{\lambda'}(i)=m_\lambda(i)$ for $i\notin\{e,e-2,e-j,e-j-1\}$, $\chi'(e-k)=f-1$ for $k\in[1,j]$, $\chi'(i)=\chi(i)$ for $i\leq e-j-1$.

(iii) $e<2f$. We have

$m_{\lambda'}(e-j)=m_\lambda(e-j)-2$, $m_{\lambda'}(e-j-1)=m_\lambda(e-j-1)+2$ (if $e>j+1$), $m_{\lambda'}(i)=m_\lambda(i)$ for $i\notin\{e-j,e-j-1\}$, $\chi'(e-k)=f-1$ for $k\in[0,j]$, $\chi'(i)=\chi(i)$ for $i\leq e-j-1$.

Let $\alpha_{\xi'}$ and $T_{\xi'}$ be defined for ξ' as before. Let $m_\lambda(e-i)=2d_i$.

Assume that we are in case (i). We have a decomposition $V=W\oplus Y$ of V into mutually orthogonal T_ξ-stable subspaces such that

$$\alpha_\xi|_W={}^*W_f(e)^{d_0}\text{ and }T_\xi^{e-1}Y=0.$$

Then $V_{\geq -N+1}=(\ker T_\xi^{e-1}\cap W)\oplus Y$ and $V_{\geq N}=T_\xi^{e-1}W$. Hence we have a natural decomposition $V'=W'\oplus Y$ of V' into mutually orthogonal $T_{\xi'}$-stable subspaces, where $W'=(\ker T_\xi^{e-1}\cap W)/T_\xi^{e-1}W$, and (see 2.4.2 (ii))

$$\alpha_{\xi'}|_{W'}={}^*W_{f-1}(e-2)^{d_0},\ \alpha_{\xi'}|_Y=\alpha_\xi|_Y.$$

We have $\chi'(i)=\max(\chi_{\alpha_{\xi'}|_{W'}}(i),\chi_{\alpha_\xi|_Y}(i))$ and $\chi(i)=\max(\chi_{\alpha_\xi|_W}(i),\chi_{\alpha_\xi|_Y}(i))$. For $0<\lambda_i\leq e-3$, $\lambda_i-\chi(\lambda_i)\leq\frac{\lambda_i+1}{2}<e-f$ and thus $\chi_{\alpha_\xi|_W}(\lambda_i)=\max(0,\lambda_i-e+f)<\chi(\lambda_i)$, which implies that $\chi_{\alpha_\xi|_Y}(\lambda_i)=\chi(\lambda_i)$ and thus $\chi'(\lambda_i)=\max(\max(\lambda_i-e+f+1,0),\chi_{\alpha_\xi|_Y}(\lambda_i))=\chi(\lambda_i)$. Now $\chi'(e-2)=\max(f-1,\chi_{\alpha_\xi|_Y}(e-2))$ and $\chi(e-2)=\max(f-2,\chi_{\alpha_\xi|_Y}(e-2))$, thus the assertion on $\chi'(e-2)$ holds; $\chi'(e-1)=\chi(e-1)$ since $\chi_{\alpha_{\xi'}|_{W'}}(e-1)=\chi_{\alpha_\xi|_W}(e-1)$.

Assume that we are in case (ii). Then $j\geq 1$. We have a decomposition $V=W_0\oplus W_1\oplus Y$ of V into mutually orthogonal T-stable subspaces such that (we use [X1, Lemma 2.9])

$$\alpha_\xi|_{W_0}={}^*W_f(e)^{d_0},\ \alpha_\xi|_{W_1}={}^*W_f(e-j),\ T_\xi^{e-1}Y=0\text{ and }\alpha_\xi(T_\xi^{f-1}Y)=0.$$

Then $V_{\geq -N+1}=(\ker T_\xi^{e-1}\cap W_0)\oplus K_{W_1}\oplus Y$ and $V_{\geq N}=T_\xi^{e-1}W_0\oplus L_{W_1}$, where $K_{W_1}=\{v\in W_1|\alpha_\xi(T_\xi^{f-1}v)=0\}$ and $L_{W_1}=K_{W_1}^\perp\cap W_1$. Hence we have a natural decomposition $V'=W_0'\oplus W_1'\oplus Y$ of V' into mutually orthogonal $T_{\xi'}$-stable subspaces, where $W_0'=(\ker T_\xi^{e-1}\cap W_0)/T_\xi^{e-1}W_0$, $W_1'=K_{W_1}/L_{W_1}$, and (see 2.4.2 (i) (ii))

$$\alpha_{\xi'}|_{W_0'}={}^*W_{f-1}(e-2)^{d_0},\ \alpha_{\xi'}|_{W_1'}={}^*W_{f-1}(e-j-1),\ \alpha_{\xi'}|_Y=\alpha_\xi|_Y.$$

We have $\chi'(i)=\max(\chi_{\alpha_{\xi'}|_{W_0'}}(i),\chi_{\alpha_{\xi'}|_{W_1'}}(i),\chi_{\alpha_\xi|_Y}(i))$ and $\chi(i)=\max(\chi_{\alpha_\xi|_{W_0}}(i),\chi_{\alpha_\xi|_{W_1}}(i),\chi_{\alpha_\xi|_Y}(i))$. Thus for $e-j\leq i\leq e-1$, $\chi'(i)=f-1$ (note that $\chi_{\alpha_\xi|_Y}(i)\leq f-1$); for $i\leq e-j-1$, $\chi_{\alpha_\xi|_{W_0}}(i)\leq\chi_{\alpha_{\xi'}|_{W_0'}}(i)\leq\chi_{\alpha_{\xi'}|_{W_1'}}(i)=\chi_{\alpha_\xi|_{W_1}}(i)$ (since $j\geq 1$) and thus $\chi'(i)=\chi(i)$.

Assume that we are in case (iii). We have a decomposition of V into mutually orthogonal T_ξ-stable subspaces $V=W\oplus Y$ such that (we use [X1, Lemma 2.9])

$$\alpha_\xi|_W={}^*W_f(e-j)\text{ and }\alpha_\xi(T_\xi^{f-1}Y)=0.$$

Then $V_{\geq -N+1}=K_W\oplus Y$ and $V_{\geq N}=L_W$, where $K_W=\{v\in W|\alpha_\xi(T_\xi^{f-1}v)=0\}$ and $L_W=K_W^\perp\cap W$. Hence we have a natural decomposition $V'=W'\oplus Y$ of V' into mutually orthogonal T'-stable subspaces, where $W'=K_W/L_W$. Moreover (see 2.4.2 (i))

$$\alpha_{\xi'}|_{W'} = {}^*W_{f-1}(e-j-1) \text{ and } \alpha_{\xi'}|_Y = \alpha_\xi|_Y.$$

We have $\chi'(i) = \max(\chi_{\alpha_{\xi'}|_{W'}}(i), \chi_{\alpha_\xi|_Y}(i))$ and $\chi(i) = \max(\chi_{\alpha_\xi|_W}(i), \chi_{\alpha_\xi|_Y}(i))$. Thus for $i \geq e-j$, $\chi'(i) = f-1$ (as $\chi_{\alpha_\xi|_Y}(i) \leq f-1$); for $i \leq e-j-1$, $\chi_{\alpha_{\xi'}|_{W'}}(i) = \chi_{\alpha_\xi|_W}(i)$ and thus $\chi'(i) = \chi(i)$.

6.6. Using the definition of Ψ^*_R and the description of c' in 6.5, we compute $\Psi^*_R(\mathrm{c}) = \tilde{\lambda}$ and $\Psi^*_{R'}(\mathrm{c}') = \tilde{\lambda}'$ in each case (i)-(iii) as follows. It is then easy to check that 6.3 (b) holds in each case.

Let $d = d_0 + d_1$ in case (i) and $d = \sum_{a\in[0,j]} d_a$ in cases (ii) and (iii). We have $\tilde{\lambda}_i = \tilde{\lambda}'_i$ for all $i \geq 2d+2$, since $\mu_i = \mu'_i$ and $\nu_i = \nu'_i$ for all $i \geq d+1$. In case (i), if $\chi(e-2) \leq f-1$, then $\tilde{\lambda}_{2d+1} = \tilde{\lambda}'_{2d+1}$ since $\mu_{d+1} \leq \nu_d$ and $\mu'_{d+1} \leq \nu'_d$. In cases (ii) and (iii), $\tilde{\lambda}_i = \tilde{\lambda}'_i$ for $i = 2d, 2d+1$, since $\nu_d = \nu'_d$, $\nu_d \leq \mu_d$ and $\nu'_d \leq \mu'_d$. Let $\tilde{\lambda}^1 = (\tilde{\lambda}_1, \ldots, \tilde{\lambda}_{2d+1})$ and $\tilde{\lambda}'^1 = (\tilde{\lambda}'_1, \ldots, \tilde{\lambda}'_{2d+1})$. We have

(i)
$$\begin{array}{ll} \tilde{\lambda}^1 = e^{2d_0}(e-1)^{2d_1+1},\ \tilde{\lambda}'^1 = (e-1)^{2d_1+1}(e-2)^{2d_0} & \text{if } \chi(e-2) = f, \\ \tilde{\lambda}^1 = e^{2d_0}(e-1)^{2d_1}\tilde{\lambda}_{2d+1},\ \tilde{\lambda}'^1 = (e-1)^{2d_1}(e-2)^{2d_0}\tilde{\lambda}_{2d+1} & \text{if } \chi(e-2) \leq f-1; \end{array}$$

(ii)
$$\begin{array}{l} \tilde{\lambda}^1 = e^{2d_0+1}(e-1)^{2d_1}(e-2)^{2d_2}\cdots(e-j+1)^{2d_{j-1}}(e-j)^{2d_j-2}\tilde{\lambda}_{2d}\tilde{\lambda}_{2d+1}, \\ \tilde{\lambda}'^1 = (e-2)^{2d_0+1}(e-1)^{2d_1}(e-2)^{2d_2}\cdots(e-j+1)^{2d_{j-1}}(e-j)^{2d_j-2}\tilde{\lambda}_{2d}\tilde{\lambda}_{2d+1}; \end{array}$$

(iii)
$$\begin{array}{l} \tilde{\lambda}^1 = (2f)e^{2d_0}(e-1)^{2d_1}(e-2)^{2d_2}\cdots(e-j+1)^{2d_{j-1}}(e-j)^{2d_j-2}\tilde{\lambda}_{2d}\tilde{\lambda}_{2d+1}, \\ \tilde{\lambda}'^1 = (2f-2)e^{2d_0}(e-1)^{2d_1}(e-2)^{2d_2}\cdots(e-j+1)^{2d_{j-1}}(e-j)^{2d_j-2}\tilde{\lambda}_{2d}\tilde{\lambda}_{2d+1}. \end{array}$$

6.7. Assume that $p = 2$ and $G = SO(V, Q)$ in the remainder of this section. Suppose that $\mathrm{c} = (m, (\lambda, \chi))$. We keep the notations in 6.3. We show in this subsection that

$$\text{(a) } f_N = \begin{cases} 1 & \textit{if } m \geq f, \\ 2 & \textit{if } e-f < m < f, \\ m_\lambda(e) & \textit{if } m = 0,\ \textit{or } 0 < m = e-f < f-1 \textit{ and } \rho = 0, \\ m_\lambda(e)+1 & \textit{if } 0 < m = e-f = f-1 \textit{ and } \rho = 0, \\ m_\lambda(e)+2 & \textit{if } 0 < m = e-f < f \textit{ and } \rho \neq 0. \end{cases}$$

Recall that $f_N = \dim V_{\geq N}$ and $V_{\geq N} = V^{\perp}_{\geq -N+1} \cap Q^{-1}(0)$ (see 2.4.3). We describe $V_{\geq N}$ in various cases in the following and then (a) follows.

Suppose that $m = 0$. Then $e \geq 2$ since $\xi \neq 0$. We have $V_{\geq N} = \{w + \sqrt{Q(w)}v_0 | w \in \mathrm{Im} T_\xi^{e-1}\}$, where $\sqrt{}$ is a chosen square root on $\mathbf{k}$.

Suppose that $m \geq f$. Then $V_{\geq N} = \mathrm{span}\{v_0\}$.

Suppose that $e - f < m < f$. Then $V_{\geq N} = \mathrm{span}\{v_0\} \oplus L$, where $L = \{w \in W | Q(T_\xi^{f-1}w) = 0\}^\perp \cap W$ is a line (see [X2, 5.5]).

Suppose that $0 < m = e-f = f-1$. Then $V_{\geq N} = \mathrm{span}\{v_0\} \oplus L$, where $L = \{w \in W | T_\xi^{e-1}w = 0, Q(T_\xi^{f-1}w) = 0\}^\perp \cap W$, and $\dim L = m_\lambda(e)$ if $\rho = 0$, $\dim L = m_\lambda(e)+1$ if $\rho \neq 0$ (see [X2, 5.5]).

Suppose that $0 < m = e-f < f-1$ and $\rho \neq 0$. Then $V_{\geq N} = \mathrm{span}\{v_0\} \oplus L$, where $L = \{w \in W | T_\xi^{e-1}w = 0, Q(T_\xi^{f-1}w) = 0\}^\perp \cap W$. Same argument as in [X2, 5.5] shows that we have a decomposition $W = W_1 \oplus W_2$ of W into T_ξ-stable orthogonal subspaces such that $T_\xi^{e-1}W_2 = 0$ and $\chi_{T_\xi|_{W_2}}(e-1) = f$. Hence $L = \mathrm{Im}T_\xi^{e-1} \oplus (\{x \in W_2 | Q(T_\xi^{f-1}x) = 0\}^\perp \cap W_2)$ and $\dim L = m_\lambda(e)+1$.

Suppose that $0 < m = e-f < f-1$ and $\rho = 0$. Then $V_{\geq N} = \{w + \beta(w, w_{**})v_0 | w \in \mathrm{Im}T_\xi^{e-1}\}$.

6.8. We describe $\mathrm{c}' = (m'; (\lambda', \chi'))$ in various cases as follows. Let $j \geq 0$ be the unique integer such that $\chi(e-j) = f$ and $\chi(e-j-1) < f$.

(i) $m = 0$, or $0 < e - f = m < f - 1$ and $\rho = 0$. We have

$m' = m$, $m_{\lambda'}(e) = 0$, $m_{\lambda'}(e-2) = m_\lambda(e-2) + m_\lambda(e)$ *(if* $e > 2$*),* $m_{\lambda'}(i) = m_\lambda(i)$ *for* $i \notin \{e, e-2\}$, $\chi'(i) = \chi(i)$ *for* $i \leq e - 1$.

(ii) $m \geq f$. We have

$m' = m - 1$, $m_{\lambda'}(i) = m_\lambda(i)$ *for all* i, $\chi'(\lambda_i) = \chi(\lambda_i)$ *for* $\lambda_i \leq e - 1$, $\chi'(e) = f + 1$ *if* $m = e - f$, *and* $\chi'(e) = f$ *if* $m > e - f$.

(iii) $e - f < m < f$. We have

$m' = m - 1, m_{\lambda'}(e-j) = m_\lambda(e-j) - 2$, $m_{\lambda'}(e-j-1) = m_\lambda(e-j-1) + 2$ *(if* $e > j+1$*),* $m_{\lambda'}(i) = m_\lambda(i)$ *for* $i \notin \{e-j, e-j-1\}$, $\chi'(\lambda_i) = f - 1$ *for* $e - j \leq \lambda_i \leq e - 1$, $\chi'(\lambda_i) = \chi(\lambda_i)$ *for* $\lambda_i \leq e - j - 1$, $\chi'(e) = f - 1$ *if* $m \geq e - f + 2$ *and* $\chi'(e) = f$ *if* $m = e - f + 1$.

(iv) $0 < e - f = m < f$ and $\rho \neq 0$. We have

$m' = m - 1$, $m_{\lambda'}(e) = 0$, $m_{\lambda'}(e-2) = m_\lambda(e-2) + m_\lambda(e) + 2\delta_{j,1} - 2\delta_{j,2}$ *(if* $e > 2$*),* $m_{\lambda'}(e-j) = m_\lambda(e-j) - 2 + \delta_{j,2} m_\lambda(e)$, $m_{\lambda'}(e-j-1) = m_\lambda(e-j-1) + 2 + \delta_{j,1} m_\lambda(e)$ *(if* $e > j + 1$*),* $m_{\lambda'}(i) = m_\lambda(i)$ *for* $i \notin \{e, e-2, e-j, e-j-1\}$, $\chi'(e-1) = f$, $\chi'(\lambda_i) = f - 1$ *for* $e - j \leq \lambda_i \leq e - 2$, $\chi'(\lambda_i) = \chi(\lambda_i)$ *for* $\lambda_i \leq e - j - 1$.

(v) $0 < e - f = m = f - 1$ and $\rho = 0$. We have

$m' = m - 1, m_{\lambda'}(e) = 0$, $m_{\lambda'}(e-2) = m_\lambda(e) + m_\lambda(e-2)$ *(if* $e > 2$*),* $m_{\lambda'}(i) = m_\lambda(i)$ *for* $i \notin \{e, e-2\}$, $\chi'(e-1) = f$, $\chi'(e-2) = f - 1$, $\chi'(\lambda_i) = \chi(\lambda_i)$ *for* $\lambda_i \leq e - 2$.

Recall that we can choose u_0 (or W if $m = 0$) such that $\chi_W(\lambda_i) = \chi(\lambda_i)$ in the decomposition $V = \mathrm{span}\{u_i, v_i\} \oplus W$. In the following u_0 or W is chosen as such. Let $m_\lambda(e-i) = 2d_i$.

Assume that we are in case (i). Suppose first that $m = 0$. Then $e = f$. There exists $w_0 \in W$ such that $\beta(w_0, w)^2 = Q(w)$ for all $w \in W$. Let $\tilde{W} = \{w + \beta(w_0, w)v_0 | w \in W\}$. Then $V_{\geq -N+1} = \mathrm{span}\{v_0\} \oplus \ker \tilde{T}_\xi^{e-1}$ and $V_{\geq N} = \mathrm{Im}\tilde{T}_\xi^{e-1}$ (note that $Q(\tilde{T}_\xi^{e-1}\tilde{W}) = 0$). The description for c' follows. Suppose now that $0 < e - f = m < f - 1$ and $\rho = 0$. Let $\tilde{u}_0 = u_0 + w_{**}$ and let $\tilde{u}_i, \tilde{W}, \tilde{T}_\xi$ be defined accordingly. Then for all $\tilde{w} \in \tilde{W}$, $Q(\tilde{T}_\xi^{f-1}\tilde{w}) = \beta(T_\xi^{e-1} w_{**}, \pi_W(\tilde{w}))^2 + Q(T_\xi^{f-1}\pi_W(\tilde{w})) = 0$. Then $V' = \mathrm{span}\{v_i, i \in [0, m], \tilde{u}_i, i \in [0, m-1]\} \oplus \tilde{W}'$, where $\tilde{W}' = (\ker \tilde{T}_\xi^{e-1} / \mathrm{Im}\tilde{T}_\xi^{e-1})$. Thus $m' = m$. Let $T_{\xi'} : \tilde{W}' \to \tilde{W}'$ be defined for $\alpha_{\xi'}$. We have $\chi_{\tilde{W}}(i) \leq f - 1$. Suppose that $\chi_{\tilde{W}}(e) = \tilde{f}$. We have a decomposition $\tilde{W} = W_1 \oplus W_2$ of $\tilde{W}$ into mutually orthogonal $\tilde{T}_\xi$-stable subspaces such that (see [X3, 5.6])

$$\tilde{T}_\xi|_{W_1} = W_{\tilde{f}}(e)^{d_0} \text{ and } T_\xi^{e-1} W_2 = 0.$$

Then $\tilde{W}' = W_1' \oplus W_2$, where $W_1' = (\ker \tilde{T}_\xi^{e-1} \cap W_1)/\tilde{T}_\xi^{e-1} W_1$ and $T_{\xi'}|_{W_1'} = W_{\tilde{f}-1}(e-2)^{d_0}$. We have $\chi'(i) = \max(i - m, \chi_{T_{\xi'}|_{W_1'}}(i), \chi_{\tilde{T}_\xi|_{W_2}}(i))$ and $\chi(i) = \max(i - m, \chi_{\tilde{T}_\xi|_{W_1}}(i), \chi_{\tilde{T}_\xi|_{W_2}}(i))$. Thus $\chi'(e-1) = f - 1 = \chi(e-1)$ and $\chi'(i) = \max(i - m, \chi_{\tilde{T}_\xi|_{W_2}}(i)) = \chi(i)$ (note that $\chi_{T_{\xi'}|_{W_1'}}(i) \leq \max(0, i-m)$ and $\chi_{T_\xi|_{W_1}}(i) \leq \max(0, i-m)$) for all $i \leq e - 2$.

Assume that we are in case (ii). Then $V' = (\mathrm{span}\{v_i, u_i, i \in [0, m-1], v_m\}/\mathrm{span}\{v_0\}) \oplus W$. Thus $m' = m - 1$ and $m_{\lambda'}(i) = m_\lambda(i)$. We have $\chi'(\lambda_i) = \max(\lambda_i - m + 1, \chi(\lambda_i))$. If $m > \lambda_i - \chi(\lambda_i)$, then $\chi'(\lambda_i) = \chi(\lambda_i)$; if $m = \lambda_i - \chi(\lambda_i)$, then $m = e - f$ and thus $e = 2m$, $\lambda_i = e$ (note that $\chi(\lambda_i) \geq \lambda_i/2 > \lambda_i - m$ for $\lambda_i < e = 2m$) and $\chi'(e) = f + 1$.

In cases (iii)-(v), we have $V' = (\mathrm{span}\{v_i, u_i, i \in [0, m-1], v_m\}/\mathrm{span}\{v_0\}) \oplus W'$, where $W' = \Lambda_W/(\Lambda_W^{\perp} \cap W)$ for some subspace $\Lambda_W \subset W$. Thus $m' = m-1$. Let $T_{\xi'} : W' \to W'$ be defined for $\alpha_{\xi'}$. We write $T_{\xi'} = (\lambda', \chi'_{W'})$. We can apply the results in [X3, 5.6] for T_ξ and $T_{\xi'}$ and then in each case the assertions on $m_{\lambda'}(i)$ follow. The assertions on χ' also hold since $\chi'(\lambda'_i) = \max(\lambda'_i - m + 1, \chi'_{W'}(i))$ (see below for the description of $\chi'_{W'}(i)$).

(iii) In this case $\Lambda_W = \{w \in W | Q(T_\xi^{f-1} w) = 0\}$ and $\chi'_{W'}(e-k) = f-1$ for $k \in [0, j]$, $\chi'_{W'}(\lambda_i) = \chi(\lambda_i)$ for $\lambda_i \leq e - j - 1$. Note that $\lambda_i - m + 1 \leq \chi(\lambda_i)$ for $\lambda_i \leq e - j - 1$.

(iv) In this case $\Lambda_W = \{w \in W | T_\xi^{e-1} w = 0, Q(T_\xi^{f-1} w) = 0\}$, and $\chi'_{W'}(e-k) = f - 1$ for $k \in [0, j]$, $\chi'_{W'}(\lambda_i) = \chi(\lambda_i)$ for $\lambda_i \leq e - j - 1$. Since $\rho \neq 0$, $\chi(e-1) = f$ and thus $m > \lambda_i - \chi(\lambda_i)$ for all $\lambda_i \leq e - 1$.

(v) In this case $\Lambda_W = \{w \in W | T_\xi^{e-1} w = 0\}$ and $\chi'_{W'}(\lambda_i) = \chi(\lambda_i)$ for $\lambda_i \leq e - 1$. Note that for $\lambda_i \leq e - 2$, $\chi(\lambda_i) \geq [\frac{\lambda_i + 1}{2}] \geq \lambda_i - m + 1$.

6.9. Using the definition of Ψ_R^* and the description of c' in 6.8, we compute $\Psi_R^*(\mathrm{c}) = \tilde{\lambda}$ and $\Psi_{R'}^*(\mathrm{c}') = \tilde{\lambda}'$ in each case (i)-(v) as follows. It is then easy to check that 6.3 (b) holds in each case.

Let $d = d_0 + d_1$ in cases (i) and (v); $d = d_0$ in case (ii); and $d = \sum_{a \in [0,j]} d_a$ in cases (iii) and (iv). Then $\tilde{\lambda}'_i = \tilde{\lambda}_i$ for all $i \geq 2d + 3$ since $\mu_i = \mu'_i$ and $\nu_{i-1} = \nu'_{i-1}$ for all $i \geq d + 2$. In case (i), if $\chi(e-2) = f - 2$, then $\mu_{d+1} = \mu'_{d+1}$ and thus $\tilde{\lambda}'_{2d+2} = \tilde{\lambda}'_{2d+2}$; if moreover $f \geq m + 3$, then $\tilde{\lambda}'_{2d+1} = \tilde{\lambda}'_{2d+1}$ since $\mu_{d+1} < \nu_d$, $\mu'_{d+1} < \nu'_d$. In case (v) or in case (ii) with $m = e - f$ (then $e = 2f$), $\tilde{\lambda}_{2d+2} = \tilde{\lambda}'_{2d+2}$ since $\nu_{d+1} < \mu_{d+1} + 2$ and $\nu'_{d+1} < \mu'_{d+1} + 2$; if $m > e - f$, then $\tilde{\lambda}_i = \tilde{\lambda}'_i$ for $i = 2d+1, 2d+2$ since $\mu_{d+1} = \mu'_{d+1}$ and $\nu_d = \nu'_d$. In case (iii) and (iv), $\tilde{\lambda}_i = \tilde{\lambda}'_i$ for $i = 2d+1, 2d+2$ since $\mu_{d+1} = \mu'_{d+1}$ and $\mu_{d+1} < \nu_d$, $\mu'_{d+1} < \nu'_d$.

Let $\tilde{\lambda}^1 = (\tilde{\lambda}_1, \ldots, \tilde{\lambda}_{2d+2})$ and $\tilde{\lambda}'^1 = (\tilde{\lambda}'_1, \ldots, \tilde{\lambda}'_{2d+2})$. We have

$$\text{(i)} \quad \begin{array}{ll} \tilde{\lambda}^1 = e^{2d_0}(e-1)^{2d_1}\tilde{\lambda}_{2d+1}\tilde{\lambda}_{2d+2}, & \\ \tilde{\lambda}'^1 = (e-1)^{2d_1}(e-2)^{2d_0}\tilde{\lambda}_{2d+1}\tilde{\lambda}_{2d+2} & \text{if } \chi(e-2) = f - 2 \geq m + 1 \\ \tilde{\lambda}^1 = e^{2d_0}(e-1)^{2d_1+1}\tilde{\lambda}_{2d+2},\ \tilde{\lambda}'^1 = (e-1)^{2d_1+1}(e-2)^{2d_0}\tilde{\lambda}_{2d+2} & \text{if } \chi(e-2) = f - 2 = m \\ \tilde{\lambda}^1 = e^{2d_0}(e-1)^{2d_1+2},\ \tilde{\lambda}'^1 = (e-1)^{2d_1+2}(e-2)^{2d_0} & \text{if } \chi(e-2) = f - 1. \end{array}$$

$$\text{(ii)} \quad \begin{array}{ll} \tilde{\lambda}^1 = (e+1)e^{2d_0}\tilde{\lambda}_{2d+2},\ \tilde{\lambda}'^1 = e^{2d_0}(e-1)\tilde{\lambda}_{2d+2} & \text{if } m = e - f \\ \tilde{\lambda}^1 = (2m+1)(2f-1)e^{2d_0-2}\tilde{\lambda}_{2d+1}\tilde{\lambda}_{2d+2}, & \\ \tilde{\lambda}'^1 = (2m-1)(2f-1)e^{2d_0-2}\tilde{\lambda}_{2d+1}\tilde{\lambda}_{2d+2} & \text{if } m > e - f \text{ and } e < 2f; \\ \tilde{\lambda}^1 = (2m+1)e^{2d_0-1}\tilde{\lambda}_{2d+1}\tilde{\lambda}_{2d+2},\ \tilde{\lambda}'^1 = (2m-1)e^{2d_0-1}\tilde{\lambda}_{2d+1}\tilde{\lambda}_{2d+2} & \text{if } m > e - f \text{ and } e = 2f. \end{array}$$

$$\text{(iii)} \quad \begin{array}{l} \tilde{\lambda}^1 = (m+f)^2 e^{2d_0} \cdots (e-j)^{2d_j-2}\tilde{\lambda}_{2d+1}\tilde{\lambda}_{2d+2}, \\ \tilde{\lambda}'^1 = (m+f-2)^2 e^{2d_0} \cdots (e-j)^{2d_j-2}\tilde{\lambda}_{2d+1}\tilde{\lambda}_{2d+2} \end{array} .$$

$$\text{(iv)} \quad \begin{array}{l} \tilde{\lambda}^1 = e^{2d_0+2}(e-1)^{2d_1}(e-2)^{2d_2} \cdots (e-j+1)^{2d_{j-1}}(e-j)^{2d_j-2}, \\ \tilde{\lambda}'^1 = (e-2)^{2d_0+2}(e-1)^{2d_1}(e-2)^{2d_2} \cdots (e-j+1)^{2d_{j-1}}(e-j)^{2d_j-2}. \end{array}$$

(v) $\tilde{\lambda}^1 = e^{2d_0+1}(e-1)^{2d_1}\tilde{\lambda}_{2d+2},\ \tilde{\lambda}'^1 = (e-1)^{2d_1}(e-2)^{2d_0+1}\tilde{\lambda}_{2d+2}$.

7. Nilpotent orbits in $\mathfrak{g}^*$ for G of exceptional type

7.1. Assume that G is of exceptional type G_2, F_4, E_6, E_7, or E_8. We have

Proposition 7.1. *The number of nilpotent G-orbits in $\mathfrak{g}^*$ is finite.*

In [K, Theorem 2.2], Kac constructs an invariant non-degenerate bilinear form on a symmetrizable Kac-Moody Lie algebra over $\mathbb{C}$. Assume G is of type G_2 and $p \neq 3$, of type F_4, E_6 or E_7 and $p \neq 2$, or of type E_8 (resp. G is of type E_6 or E_7 and $p = 2$). Then the Cartan matrix of $\mathfrak{g}$ is symmetrizable and one can apply the method used in [*loc.cit*] to construct an invariant non-degenerate bilinear form (resp. pairing) on $\mathfrak{g}$ (resp. between the Lie algebra $\mathfrak{g}_s$ of the simply connected group and the Lie algebra $\mathfrak{g}_{ad}$ of the adjoint group). Thus in the above cases $\mathfrak{g}^*$ and $\mathfrak{g}$ (resp. $\mathfrak{g}_s^*$ and $\mathfrak{g}_{ad}$) can be identified via the bilinear form (resp. pairing); and then the proposition follows from the corresponding result for $\mathfrak{g}$ since the nilpotent orbits in $\mathfrak{g}^*$ and $\mathfrak{g}$ (resp. in $\mathfrak{g}_s^*$ and $\mathfrak{g}_{ad}$) are identified. Hence it remains to study the nilpotent orbits in $\mathfrak{g}^*$ when G is of type G_2 and $p = 3$, or G is of type F_4 and $p = 2$. (The above argument was suggested by G. Lusztig.) The proof of the proposition in these remaining cases will be given in Subsections 7.2-7.5.

7.2. Let $\mathbf{F}_q$ be a finite field of characteristic 3 (resp. 2). Let G be of type G_2 (resp. F_4) defined over $\mathbf{F}_q$. We study the nilpotent $G(\mathbf{F}_q)$-orbits in $\mathfrak{g}(\mathbf{F}_q)^*$. The strategy is as follows. We find representatives $\xi \in \mathcal{N}_{\mathfrak{g}(\mathbf{F}_q)^*}$ for the nilpotent orbits and compute $|Z_G(\xi)(\mathbf{F}_q)|$ (the number of rational points in the centralizers $Z_G(\xi)$). Then by a direct computation one sees that the numbers of rational points in all nilpotent orbits add up to q^{2N}, where N is the number of positive roots. As the number of nilpotent elements in $\mathfrak{g}(\mathbf{F_q})^*$ equals q^{2N} (shown by G. Lusztig), we get all the nilpotent orbits.

7.3. We describe how to compute $|Z_G(\xi)(\mathbf{F}_q)|$ for $\xi \in \mathcal{N}_{\mathfrak{g}(\mathbf{F}_q)^*}$ in this subsection.

Let T be a maximal torus of G, R the root system of (G, T), Π a set of simple roots in R and R^+ the corresponding set of positive roots in R. We have a Chevalley basis $\{h_\alpha,\ \alpha \in \Pi;\ e_\alpha,\ \alpha \in R\}$ of $\mathfrak{g}$ satisfying

$$[h_\alpha, h_\beta] = 0;\ [h_\alpha, e_\beta] = A_{\alpha\beta} e_\beta;\ [e_\alpha, e_{-\alpha}] = h_\alpha;\ [e_\alpha, e_\beta] = N_{\alpha,\beta} e_{\alpha+\beta},\ \text{if } \alpha + \beta \in R,$$

where $A_{\alpha,\beta}$ and $N_{\alpha,\beta}$ are constant integers (for determination of structural constants $N_{\alpha,\beta}$ see [C]). For each $\alpha \in R$, there is a unique 1-dimensional connected closed unipotent subgroup $U_\alpha \subset G$ and an isomorphism

$$x_\alpha :\ \mathbb{G}_a \to U_\alpha$$

such that $s x_\alpha(t) s^{-1} = x_\alpha(\alpha(s)t)$ for all $s \in T$ and $t \in \mathbb{G}_a$. We assume that $dx_\alpha(1) = e_\alpha$ and $n_\alpha(t) = x_\alpha(t) x_{-\alpha}(-t^{-1}) x_\alpha(t)$ normalizes T. Define $h_\alpha(t) = n_\alpha(t) n_\alpha(-1)$. Then T is generated by $h_\alpha(\lambda), \alpha \in \Pi, \lambda \in \mathbf{k}^\times$. Let B be the Borel subgroup UT of G, where $U = \{ \prod_{\alpha \in R^+} x_\alpha(t_\alpha), t_\alpha \in \mathbb{G}_a \}$. By Bruhat decomposition, every element g in G can be written uniquely in the form

$$g = b n_w u_w$$

for some $w \in W$, some $b \in B$ and some $u_w \in U_w = \{ \prod_{\alpha > 0, w(\alpha) < 0} x_\alpha(t_\alpha) \,|\, t_\alpha \in \mathbb{G}_a \}$, where n_w is a representative of w in $N_G(T)$. We can choose $n_\alpha = n_\alpha(1)$ to be the representative of the simple reflections $s_\alpha \in W$, $\alpha \in \Pi$. Let $\mathfrak{t}$ be the Lie algebra of T and let $\mathfrak{t}^* = \{\xi \in \mathfrak{g}^* \,|\, \xi(e_\alpha) = 0,\ \forall\ \alpha \in R\}$. We define $e'_\alpha \in \mathfrak{g}^*$ by

$$e'_\alpha(\mathfrak{t}) = 0;\ e'_\alpha(e_\beta) = \delta_{-\alpha,\beta}, \forall\ \beta \in R.$$

Then $\{e'_\alpha, \alpha \in R^+\}$ form a basis of $\mathfrak{n}^*$. The coadjoint action of $x_\alpha(t)$ on e'_β, $\alpha, \beta \in R^+$ are given as follows

$$\text{(a)} \quad x_\alpha(t).e'_\beta = \sum_i (-1)^i t^i M_{\alpha,-i\alpha-\beta,i} e'_{i\alpha+\beta},$$

where $M_{\alpha,\beta,i} = \frac{1}{i!} N_{\alpha,\beta} N_{\alpha,\alpha+\beta} \cdots N_{\alpha,(i-1)\alpha+\beta}$.

We can find representatives of nilpotent orbits in $\mathfrak{n}^*$, namely, we can assume that $\xi = \sum_{\alpha \in R^+} a_\alpha e'_\alpha$ for some $a_\alpha \in \mathbb{G}_a$. Now we can compute $|Z_G(\xi)(\mathbf{F}_q)|$ using the Bruhat decomposition and the formula (a). In particular, we need knowledge on the set $\{w \in W \mid Z_G(\xi) \cap (BwB) \neq \emptyset\}$. Let Δ^ξ_{min} be the set of minimal elements in the set $\{\alpha \in R^+ \mid a_\alpha \neq 0\}$ under the order relations on R^+ ($\alpha > \beta$ if $\alpha - \beta$ can be written as a sum of positive roots). If $Z_G(\xi) \cap BwB \neq \emptyset$, then for any $\alpha \in \Delta^\xi_{\text{min}}$, there exists $\beta \in \Delta^\xi_{\text{min}}$ such that $w(\alpha) \geq \beta$.

7.4. Suppose that G is of type G_2 and $p = 3$ in this subsection. We denote by α (resp. β) the short (resp. long) simple root. The structural constants can be chosen as follows

$$N_{\alpha,\beta} = 1,\ N_{\alpha,\alpha+\beta} = 2,\ N_{\alpha,2\alpha+\beta} = 3,\ N_{\beta,3\alpha+\beta} = -1,\ N_{\alpha+\beta,2\alpha+\beta} = 3.$$

Fix $\zeta \in \mathbf{F_q} \backslash \{x^2 | x \in \mathbf{F_q}\}$ and $\varpi \in \mathbf{F_q} \backslash \{x^3 + x | x \in \mathbf{F_q}\}$. The representatives ξ for nilpotent orbits over $\mathbf{F}_q$ and $|Z_G(\xi)(\mathbf{F_q})|$ are listed in Table 1.

Orbit	Representative ξ	$\lvert Z_G(\xi)(\mathbf{F_q})\rvert$
G_2	$\xi_1 = e'_\alpha + e'_\beta$	$\mathbf{q}^2$
$G_2(a_1)$	$\xi_{2,1} = e'_\beta + e'_{2\alpha+\beta}$	$6\mathbf{q}^4$
$G_2(a_1)$	$\xi_{2,2} = e'_\beta + e'_{2\alpha+\beta} - \varpi e'_{3\alpha+\beta}$	$3\mathbf{q}^4$
$G_2(a_1)$	$\xi_{2,3} = e'_\beta - \zeta e'_{2\alpha+\beta}$	$2\mathbf{q}^4$
$\widetilde{A_1}$	$\xi_3 = e'_\alpha$	$\mathbf{q}^4(\mathbf{q}^2 - 1)$
A_1	$\xi_4 = e'_\beta$	$\mathbf{q}^6(\mathbf{q}^2 - 1)$
$\emptyset$	$\xi_5 = 0$	$\mathbf{q}^6(\mathbf{q}^2 - 1)(\mathbf{q}^6 - 1)$

TABLE 1.

One can easily verify that $\xi_{2,1}$, $\xi_{2,2}$, and $\xi_{2,3}$ are in the same G-orbit. Thus $\xi_1, \xi_{2,1}, \xi_3, \xi_4, \xi_5$ form a set of representatives for nilpotent G-obits in $\mathfrak{g}^*$, and Proposition 7.1 follows.

The order relations on nilpotent orbits in $\mathfrak{g}^*$ are as follows.



7.5. Suppose that G is of type F_4 and $p=2$ in this subsection. We denote by p,q (resp. r,s) the long (resp. short) simple roots with $(q,r)\neq 0$. We denote by $apbqcrds$ the root $ap+bq+cr+ds$. The structural constants can be chosen as follows:

$$\begin{aligned}
&N_{p,q}=N_{p,qr}=N_{p,q2r}=N_{p,p3q4r2s}=N_{p,qrs}=N_{p,q2rs}=N_{p,q2r2s}=N_{q,rs}=N_{q,pq2r}=N_{q,pq2rs}\\
&\quad=N_{q,pq2r2s}=N_{q,r}=N_{q,p2q4r2s}=N_{r,s}=N_{r,qrs}=N_{r,pqrs}=N_{r,p2q2rs}=N_{pq,rs}=N_{pq,p2q4r2s}\\
&\quad=N_{s,q2r}=N_{s,pq2r}=N_{s,p2q2r}=N_{rs,pqr}=N_{rs,p2q2r}=N_{qr,pqrs}=N_{qr,pq2r2s}=N_{q2r,pq2r2s}\\
&\quad=N_{q2r,p2q2r2s}=N_{pq2r,p2q2r2s}=N_{pqr,q2rs}=N_{qrs,pq2rs}=1,\\
&N_{r,pq}=N_{r,p2q2r2s}=N_{s,qr}=N_{s,pqr}=N_{s,p2q3rs}=N_{pq,q2r}=N_{pq,q2rs}=N_{pq,q2r2s}=N_{qr,rs}\\
&\quad=N_{qr,pq2rs}=N_{rs,p2q2rs}=N_{q2r,pqrs}=N_{pqr,q2r2s}=N_{pqr,qrs}=N_{pq2r,q2r2s}=N_{qrs,pq2r}\\
&\quad=N_{pqrs,q2rs}=N_{p2q2r,pq2r2s}=N_{p2q2r,q2r2s}=-1,\\
&N_{r,pqr}=N_{r,qr}=N_{r,p2q3r2s}=N_{qr,p2q3r2s}=N_{qr,pqr}=N_{rs,pqrs}=N_{rs,qrs}\\
&\quad=N_{pqr,p2q3r2s}=N_{qrs,pqrs}=N_{q2rs,pq2rs}=N_{q2rs,p2q2rs}=N_{pq2rs,p2q2rs}=-2,\\
&N_{s,pq2rs}=N_{s,p2q2rs}=N_{s,q2rs}=N_{rs,p2q3rs}=N_{qrs,p2q3rs}=N_{pqrs,p2q3rs}=2.
\end{aligned}$$

Fix $\eta\in\mathbf{F_q}\backslash\{x^2+x|x\in\mathbf{F_q}\}$ and $\varpi\in\mathbf{F_q}\backslash\{x^3+x|x\in\mathbf{F_q}\}$. The representatives ξ for nilpotent orbits over $\mathbf{F}_q$ and $|Z_G(\xi)(\mathbf{F_q})|$ are listed in Table 2.

Orbit	Representative ξ	$\lvert Z_G(\xi)(\mathbf{F_q})\rvert$
F_4	$\xi_1=e'_p+e'_q+e'_r+e'_s$	$\mathbf{q}^4$
$F_4(a_1)$	$\xi_{2,1}=e'_p+e'_{qr}+e'_{q2r}+e'_s$	$2\mathbf{q}^6$
$F_4(a_1)$	$\xi_{2,2}=e'_p+e'_q+e'_{qr}+e'_s+\eta e'_{q2r}$	$2\mathbf{q}^6$
$F_4(a_2)$	$\xi_3=e'_p+e'_{qr}+e'_{rs}+e'_{q2r2s}$	$\mathbf{q}^8$
B_3	$\xi_4=e'_p+e'_{qrs}+e'_{q2r}+e'_{pq2rs}$	$\mathbf{q}^{10}$
C_3	$\xi_5=e'_s+e'_{q2r}+e'_{pqr}$	$\mathbf{q}^8(\mathbf{q}^2-1)$
$F_4(a_3)$	$\xi_{6,1}=e'_{pqr}+e'_{qrs}+e'_{pq2r}+e'_{q2r2s}$	$24\mathbf{q}^{12}$
$F_4(a_3)$	$\xi_{6,2}=e'_{pq}+e'_{pqr}+e'_{q2rs}+e'_{q2r2s}+\eta e'_{pq2r}$	$8\mathbf{q}^{12}$
$F_4(a_3)$	$\xi_{6,3}=e'_{pqr}+e'_{qrs}+e'_{pq2r}+e'_{q2r2s}+\eta e'_{pq2r2s}$	$4\mathbf{q}^{12}$
$F_4(a_3)$	$\xi_{6,4}=e'_{pq}+e'_{pqr}+e'_{q2rs}+e'_{q2r2s}+\eta e'_q$	$4\mathbf{q}^{12}$
$F_4(a_3)$	$\xi_{6,5}=e'_{pqr}+e'_{qrs}+e'_{q2r}+e'_{q2r2s}+\varpi e'_{pq2r2s}$	$3\mathbf{q}^{12}$
$(B_3)_2$	$\xi_7=e'_p+e'_{qr}+e'_{q2r2s}$	$\mathbf{q}^{10}(\mathbf{q}^2-1)$
$C_3(a_1)$	$\xi_{8,1}=e'_{pqr}+e'_{q2rs}+e'_{q2r2s}$	$2\mathbf{q}^{12}(\mathbf{q}^2-1)$
$C_3(a_1)$	$\xi_{8,2}=e'_{pq}+e'_{pqr}+e'_{q2rs}+\eta e'_{pq2r}$	$2\mathbf{q}^{12}(\mathbf{q}^2-1)$
B_2	$\xi_{9,1}=e'_{pqr}+e'_{q2r2s}$	$2\mathbf{q}^{12}(\mathbf{q}^2-1)^2$
B_2	$\xi_{9,2}=e'_{pq}+e'_{pqr}+e'_{q2r2s}+\eta e'_{pq2r}$	$2\mathbf{q}^{12}(\mathbf{q}^4-1)$
$\widetilde{A_2}+A_1$	$\xi_{10}=e'_{pqr}+e'_{q2rs}+e'_{p2q2r2s}$	$\mathbf{q}^{14}(\mathbf{q}^2-1)$
$A_2+\widetilde{A_1}$	$\xi_{11}=e'_{p2q2r}+e'_{q2r2s}+e'_{pq2rs}$	$\mathbf{q}^{16}(\mathbf{q}^2-1)$
$\widetilde{A_2}$	$\xi_{12}=e'_{pqrs}+e'_{q2rs}$	$\mathbf{q}^{14}(\mathbf{q}^2-1)(\mathbf{q}^6-1)$
A_2	$\xi_{13}=e'_{p2q2r}+e'_{pq2r2s}+e'_{p2q3r2s}$	$\mathbf{q}^{20}(\mathbf{q}^2-1)$
$A_1+\widetilde{A_1}$	$\xi_{14}=e'_{p2q2r2s}+e'_{p2q3rs}$	$\mathbf{q}^{20}(\mathbf{q}^2-1)^2$
$(A_2)_2$	$\xi_{15}=e'_{p2q2r}+e'_{pq2r2s}$	$\mathbf{q}^{20}(\mathbf{q}^2-1)(\mathbf{q}^6-1)$
$\widetilde{A_1}$	$\xi_{16,1}=e'_{p2q3r2s}$	$2\mathbf{q}^{21}(\mathbf{q}^2-1)(\mathbf{q}^3-1)(\mathbf{q}^4-1)$
$\widetilde{A_1}$	$\xi_{16,2}=e'_{p2q2r2s}+e'_{p2q3r2s}+\eta e'_{p2q4r2s}$	$2\mathbf{q}^{21}(\mathbf{q}^2-1)(\mathbf{q}^3+1)(\mathbf{q}^4-1)$
A_1	$\xi_{17}=e'_{2p3q4r2s}$	$\mathbf{q}^{24}(\mathbf{q}^2-1)(\mathbf{q}^4-1)(\mathbf{q}^6-1)$
$\emptyset$	$\xi_{18}=0$	$\mathbf{q}^{24}(\mathbf{q}^2-1)(\mathbf{q}^6-1)(\mathbf{q}^8-1)(\mathbf{q}^{12}-1)$

TABLE 2.

One can easily verify that $\{\xi_{2,1},\xi_{2,2}\}$, $\{\xi_{6,i}, i=1,\dots,5\}$, $\{\xi_{8,1},\xi_{8,2}\}$, $\{\xi_{9,1},\xi_{9,2}\}$, $\{\xi_{16,1},\xi_{16,2}\}$ are in the same G-orbit respectively; and ξ_i, $i=1,3,4,5,7,10,\dots,15,17,18$, $\xi_{j,1}$, $j=2,6,8,9,16$, form a set of representatives for nilpotent G-obits in $\mathfrak{g}^*$. Hence Proposition 7.1 follows.

The oder relations on nilpotent orbits in $\mathfrak{g}^*$ are as follows.



References

[C] C. Chevalley, Sur certains groupes simples. (French) Tôhoku Math. J. (2) 7 (1955), 14-66.
[H] W. H. Hesselink, Nilpotency in classical groups over a field of characteristic 2. Math. Z. 166 (1979), no. 2, 165-181.
[K] V. Kac, Infinite-dimensional Lie algebras. Third edition. Cambridge University Press, Cambridge, 1990.
[KW] V. Kac and B. Weisfeiler, Coadjoint action of a semi-simple algebraic group and the center of the enveloping algebra in characteristic p. Nederl. Akad. Wetensch. Proc. Ser. A 79=Indag. Math. 38 (1976), no. 2, 136-151.
[L1] G. Lusztig, A class of irreducible representations of a Weyl group. Nederl. Akad. Wetensch. Indag. Math. 41 (1979), no. 3, 323-335.
[L2] G. Lusztig, Intersection cohomology complexes on a reductive group. Invent. Math. 75 (1984), no.2, 205-272.
[L3] G. Lusztig, Notes on unipotent classes. Asian J. Math. 1 (1997), no. 1, 194-207.
[L4] G. Lusztig, Unipotent classes and special Weyl group representations. J. Algebra 321 (2009), no. 11, 3418-3449.
[L5] G. Lusztig, Unipotent elements in small characteristic, IV. Transform. Groups 15 (2010), no. 4, 921-936.
[LS] G. Lusztig and N. Spaltenstein, Induced unipotent classes. J. London Math. Soc. (2) 19 (1979), no. 1, 41-52.
[S1] N. Spaltenstein, On the fixed point set of a unipotent element on the variety of Borel subgroups. Topology 16 (1977), no. 2, 203-204.

[S2] N. Spaltenstein, Nilpotent Classes and Sheets of Lie Algebras in Bad Characteristic. Math. Z. 181 (1982), 31-48.

[S3] N. Spaltenstein, On the generalized Springer correspondence for exceptional groups. Algebraic groups and related topics (Kyoto/Nagoya, 1983), 317-338, Adv. Stud. Pure Math., 6, North-Holland, Amsterdam, 1985.

[St] R. Steinberg, Conjugacy classes in algebraic groups. Lecture Notes in Mathematics, Vol. 366. Springer-Verlag, Berlin-New York, 1974.

[X1] T. Xue, Nilpotent orbits in the dual of classical Lie algebras in characteristic 2 and the Springer correspondence. Represent. Theory 13 (2009), 609-635 (electronic).

[X2] T. Xue, On unipotent and nilpotent pieces for classical groups. J. Algebra 349 (2012), no. 1, 165-184.

[X3] T. Xue, Combinatorics of the Springer correspondence for classical Lie algebras and their duals in characteristic 2. Adv. Math. 230 (2012) 229-262.

[X4] T. Xue, Nilpotent elements in the dual of odd orthogonal algebras. Transform. Groups. 17 (2012), no. 2 (2012), 571-592.

Department of Mathematics, Northwestern University, Evanston, IL 60208, USA

E-mail address: `txue@math.northwestern.edu`